

5-28-00

Box 454

08/25/00
JC913 U.S. PTOJC586 U.S. PTO
09/648569
08/25/00

STANT COMMISSIONER FOR PATENTS, BOX PATENT APPLICATION, Washington, D.C. 20231

Submitted herewith for filing is the

- ☒ patent application of
☐ continuation application of
☐ continuation-in-part patent application of
- ☐ divisional application of
☐ prior application information: Examiner _____
☐ _____

Inventor(s)/Applicant Identifier: Anders, Hjelholt Pedersen et al.

For: NEW INTERFERON β -LIKE MOLECULES

- ☐ For CONTINUATION or DIVISIONAL APPS only: The entire disclosure of the prior application, from which an oath or declaration is supplied is considered a part of the disclosure of the accompanying continuation or divisional application and is hereby incorporated by reference. The incorporation can be relied upon when a portion has been inadvertently omitted from the submitted application parts.
- ☒ This application claims priority from each of the following Application Nos./filing dates: Denmark Patent Application No. PA 1999 01197, filed August 27, 1999; USSN 60/160,782, filed October 21, 1999; Denmark Patent Application No. PA 1999 01691, filed November 26, 1999; USSN 60/169,077, filed December 6, 1999; Denmark Patent Application No. PA 2000 00194, filed February 7, 2000; Denmark Patent Application No. PA 2000 00363, filed March 7, 2000; USSN 60/189,599, filed March 15, 2000; Denmark Patent Application No. PA 2000 00642, filed April 14, 2000; and USSN 60/202,248, filed May 5, 2000, the disclosure(s) of which is (are) incorporated by reference.
- ☐ Please amend this application by adding the following before the first sentence: "This application is a ☐ continuation ☐ continuation-in-part of and claims the benefit of U.S. Application No. _____, the disclosure of which is incorporated by reference for all purposes."

Enclosed are:

- ☒ 110 pages of the description (including specification, claims and abstract)
☒ 2 sheet(s) of ☐ formal ☒ informal drawing(s)
☒ 1 abstract
☒ 72 number of claims
☐ an assignment of the invention to _____
☒ a ☐ signed ☒ unsigned Declaration
☐ a Power of Attorney
☐ a certified copy of a _____ application
☒ cover sheet
☐ Information Disclosure Statement under 37 CFR 1.97
☒ Application Data Entry Sheet
- ☐ a petition to extend time to respond in the parent application
☐ notification of change of ☐ power of attorney ☐ correspondence address filed in prior application
☒ Patent Application Filing Acknowledgement postcard.
☒ 1 extra copy of this sheet is enclosed
☐ a verified statement to establish small entity status under 37 CFR 1.9 and 37 CFR 1.27
 ☐ is enclosed
 ☐ was filed in the prior application and small entity status is still proper and desired
☒ Sequence listing diskette and Statement

In view of the Unsigned Declaration as filed with this application and pursuant to 37 CFR §1.53(f),
 Applicant requests deferral of the filing fee until submission of the Missing Parts of Application.

DO NOT CHARGE THE FILING FEE AT THIS TIME.*Gwynedd Warren*

Gwynedd Warren,

Reg No.: 45,200

Patent Agent for Applicant

CORRESPONDENCE ADDRESS:

LAW OFFICES OF JONATHAN ALAN QUINE

P.O. Box 458

Alameda, CA 94501

Telephone: (510) 337-7871

Fax: (510) 337-7877

Customer No. 22798



22798

PATENT TRADEMARK OFFICE

Attorney Docket No. 02-107600US

Express Mail Label No. EL716083293US

Date of Deposit: August 25, 2000

I hereby certify that this is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above, addressed to: Assistant Commissioner for Patents Washington, Box Patent Application, D.C. 20231

By: *Tracie Brooks*

Tracie Brooks

Inventor Information

Inventor One Given Name::	Anders Hjelholt
Family Name::	Pedersen
Postal Address Line One::	58, Nybrovaenget
City::	Lyngby
State or Province::	
Country::	Denmark
Postal or Zip Code::	DK-2800
Citizenship Country::	DK
Inventor Two Given Name::	Hans Thalsgaard
Family Name::	Schambye
Postal Address Line One::	21, Pelargonievej
City::	Frederiksberg
State or Province::	
Country::	Denmark
Postal or Zip Code::	DK-2000
Citizenship Country::	DK
Inventor Three Given Name::	Kim Vilbour
Family Name::	Andersen
Postal Address Line One::	31, Taasingegade
City::	Copenhagen
State or Province::	
Country::	Denmark
Postal or Zip Code::	DK-2100
Citizenship Country::	DK
Inventor Four Given Name::	Claus
Family Name::	Bornaes
Postal Address Line One::	6, Hoejgaards Alle
City::	Hellerup
State or Province::	
Country::	Denmark
Postal or Zip Code::	DK-2900
Citizenship Country::	DK
Inventor Five Given Name::	Paul Baad
Family Name::	Rasmussen
Postal Address Line One::	4, Borthigsgade DK-2100
City::	Copenhagen
State or Province::	
Country::	Denmark
Postal or Zip Code::	DK-2100
Citizenship Country::	DK

Correspondence Information

Correspondence Customer Number:: 22798
E-mail:: jaquine@quinelaw.com

Application Information

Title Line One:: NEW INTERFERON β -LIKE MOLECULES
Title Line Two::
Total Drawing Sheets:: 2
Formal Drawings?:: No
Application Type:: Utility
Docket Number:: 02-107600US

Representative Information

Representative Customer Number:: 22798

Continuity Information

This application is a:: Non-provisional of
> Application One:: 60/160,782
Filing Date:: October 21, 1999
Patent Number::

and a :: Non-provisional of
>> Application Two:: 60/169,077
Filing Date:: December 6, 1999
Patent Number::

and a:: Non-provisional of
>>> Application Three:: 60/189,599
Filing Date:: March 15, 2000
Patent Number::

and a:: Non-provisional of
> Application Four:: 60/202,248
Filing Date:: May 5, 2000
Patent Number::

Prior Foreign Applications

Foreign Application One:: PA 1999 01197
Filing Date:: August 27, 1999
Country:: Denmark
Priority Claimed:: Yes

PATENT APPLICATION

NEW INTERFERON β -LIKE MOLECULES

Inventor(s):

Anders Hjelholt Pedersen, a citizen of Denmark, residing at:
58, Nybrovaenget, DK-2800 Lyngby DENMARK

Hans Thalsgaard Schambye, a citizen of Denmark, residing at:
21, Pelargonievej DK-2000 Frederiksberg DENMARK

Kim Vilbour Andersen, a citizen of Denmark, residing at:
31, Taasingegade DK-2100 Copenhagen DENMARK

Claus Bornaes, a citizen of Denmark, residing at:
6, Hoejgaards Alle DK-2900 Hellerup DENMARK

Paul Baad Rasmussen, a citizen of Denmark, residing at:
4, Borthigsgade DK-2100 Copenhagen DENMARK

Assignee: Maxygen ApS
1, Agern Alle, DK-2970 Hoersholm, Denmark

Entity: Large

THE LAW OFFICES OF JONATHAN ALAN QUINE

P.O. Box 458
Alameda, CA 94501
Internet address: www.quinelaw.com

Phone: (510) 337-7871
Fax: (510) 337-7877
E-mail: jaquine@quinelaw.com

NEW INTERFERON β -LIKE MOLECULES

COPYRIGHT NOTIFICATION PURSUANT TO 37 C.F.R. § 1.71(e)

A portion of the disclosure of this patent document contains material which is subject to copyright protection. The copyright owner has no objection to the facsimile reproduction by anyone of the patent document or the patent disclosure, as it appears in the Patent and Trademark Office patent file or records, but otherwise reserves all copyright rights whatsoever.

CROSS-REFERENCES TO RELATED APPLICATIONS

Pursuant to 35 USC 119 and 120, and any other applicable statute or rule, this application claims the benefit of and priority from each of the following Application Numbers/filing dates: Denmark Patent Application No. PA 1999 01197, filed August 27, 1999; USSN 60/160,782, filed October 21, 1999; Denmark Patent Application No. PA 1999 01691, filed November 26, 1999; USSN 60/169,077, filed December 6, 1999; Denmark Patent Application No. PA 2000 00194, filed February 7, 2000; Denmark Patent Application No. PA 2000 00363, filed March 7, 2000; USSN 60/189,599, filed March 15, 2000; Denmark Patent Application No. PA 2000 00642, filed April 14, 2000; and USSN 60/202,248, filed May 5, 2000, the disclosures of which are incorporated by reference.

BACKGROUND OF THE INVENTION

Interferons are important cytokines characterized by antiviral, antiproliferative, and immunomodulatory activities. These activities form a basis for the clinical benefits that have been observed in a number of diseases, including hepatitis, various cancers and multiple sclerosis. The interferons are divided into the type I and type II classes. Interferon β belongs to the class of type I interferons, which also includes interferons α , τ and ω , whereas interferon γ is the only known member of the distinct type II class.

Human interferon β is a regulatory polypeptide with a molecular weight of 22 kDa consisting of 166 amino acid residues. It can be produced by most cells in the body, in particular fibroblasts, in response to viral infection or exposure to other biologics. It binds to a multimeric cell surface receptor, and productive receptor binding results in a cascade of

5 intracellular events leading to the expression of interferon β inducible genes which in turn produces effects which can be classified as antiviral, antiproliferative and immunomodulatory.

The amino acid sequence of human interferon β was reported by Taniguchi, Gene 10:11-15, 1980, and in EP 83069, EP 41313 and US 4686191.

Crystal structures have been reported for human and murine interferon β ,
10 respectively (Proc. Natl. Acad. Sci. USA 94:11813-11818, 1997. J. Mol. Biol. 253:187-207, 1995). They have been reviewed in Cell Mol. Life Sci. 54:1203-1206, 1998.

Relatively few protein-engineered variants of interferon β have been reported (WO 9525170, WO 9848018, US 5545723, US 4914033, EP 260350, US 4588585, US 4769233, Stewart et al, DNA Vol 6 no2 1987 pp. 119-128, Runkel et al, 1998, Jour. Biol. Chem. 273, No.
15 14, pp. 8003-8008).

Expression of interferon β in CHO cells has been reported (US 4966843, US 5376567 and US 5795779).

Redlich et al, Proc. Natl. Acad. Sci., USA, Vol. 88, pp. 4040-4044, 1991 disclose immunoreactivity of antibodies against synthetic peptides corresponding to peptide stretches of
20 recombinant human interferon β with the mutation C17S.

Interferon β molecules with a particular glycosylation pattern and methods for their preparation have been reported (EP 287075 and EP 539300).

Various references disclose modification of polypeptides by polymer conjugation or glycosylation. Polymer modification of native interferon β or a C17S variant thereof has been
25 reported (EP 229108, US 5382657, EP 593868, US 4917888 and WO 99/55377). US 4,904,584 discloses PEGylated lysine depleted polypeptides, wherein at least one lysine residue has been deleted or replaced with any other amino acid residue. WO 99/67291 discloses a process for conjugating a protein with PEG, wherein at least one amino acid residue on the protein is deleted and the protein is contacted with PEG under conditions sufficient to achieve conjugation to the
30 protein. WO 99/03887 discloses PEGylated variants of polypeptides belonging to the growth hormone superfamily, wherein a cysteine residue has been substituted with a non-essential amino acid residue located in a specified region of the polypeptide. Interferon β is mentioned as one example of a polypeptide belonging to the growth hormone superfamily. WO 00/23114 discloses glycosylated and pegylated interferon β . WO 00/23472 discloses interferon β fusion

5 proteins. WO 00/26354 discloses a method of producing a glycosylated polypeptide variant with reduced allergenicity, which as compared to a corresponding parent polypeptide comprises at least one additional glycosylation site. US 5,218,092 discloses modification of granulocyte colony stimulating factor (G-CSF) and other polypeptides so as to introduce at least one additional carbohydrate chain as compared to the native polypeptide. Interferon β is mentioned
10 as one example among many polypeptides which allegedly can be modified according to the technology described in US 5,218,092.

Commercial preparations of interferon β are sold under the names Betaseron® (also termed interferon β 1b, which is non-glycosylated, produced using recombinant bacterial cells, has a deletion of the N-terminal methionine residue and the C17S mutation), and
15 Avonex™ and Rebif® (also termed interferon β 1a, which is glycosylated, produced using recombinant mammalian cells) for treatment of patients with multiple sclerosis, and have been shown to be effective in reducing the exacerbation rate. More patients treated with these interferon β agents remain exacerbation-free for prolonged periods of time as compared with placebo-treated patients. Furthermore, the accumulation rate of disability is reduced (Neurol.
20 51:682-689, 1998).

Comparison of interferon β 1a and β 1b with respect to structure and function has been presented in Pharmaceut. Res. 15:641-649, 1998.

Interferon β is the first therapeutic intervention shown to delay the progression of multiple sclerosis, a relapsing then progressive inflammatory degenerative disease of the central
25 nervous system.

Its mechanism of action, however, remains largely unclear. It appears that interferon β has inhibitory effects on the proliferation of leukocytes and antigen presentation. Furthermore, interferon β may modulate the profile of cytokine production towards an anti-inflammatory phenotype. Finally, interferon β can reduce T-cell migration by inhibiting the
30 activity of T-cell matrix metalloproteases. These activities are likely to act in concert to account for the mechanism of interferon β in MS (Neurol. 51:682-689, 1998).

In addition, interferon β may be used for the treatment of osteosarcoma, basal cell carcinoma, cervical dysplasia, glioma, acute myeloid leukemia, multiple myeloma, Hodgkin's disease, breast carcinoma, melanoma, and viral infections such as papilloma virus, viral hepatitis,

herpes genitalis, herpes zoster, herpetic keratitis, herpes simplex, viral encephalitis, cytomegalovirus pneumonia, and rhinovirus.

Various side effects are associated with the use of current preparations of interferon β , including injection site reactions, fever, chills, myalgias, arthralgias, and other flu-like symptoms (Clin. Therapeutics, 19:883-893, 1997).

In addition, 6-40% of patients develop neutralizing antibodies to interferon β (Int. Arch. Allergy Immunol. 118:368-371, 1999). It has been shown that development of interferon β -neutralizing antibodies decreases the biological response to interferon β , and cause a trend towards decreased treatment efficacy (Neurol. 50:1266-1272, 1998). Neutralizing antibodies will likely also impede the therapeutic utility of interferon β in connection with treatment of other diseases (Immunol. Immuth. 39:263-268, 1994).

Given the magnitude of side effects with current interferon β products, their association with frequent injection, the risk of developing neutralizing antibodies impeding the desired therapeutic effect of interferon β , and the potential for obtaining more optimal therapeutic interferon β levels with concomitant enhanced therapeutic effect, there is clearly a need for improved interferon β -like molecules.

SUMMARY OF THE INVENTION

This application discloses improved interferon β molecules providing one or more of the aforementioned desired benefits. In particular conjugates are disclosed that exhibit interferon β activity and comprise at least one non-polypeptide moiety covalently attached to an interferon β polypeptide that comprises an amino acid sequence that differs from that of wildtype human interferon β with the amino acid sequence shown in SEQ ID NO 2 in at least one amino acid residue selected from an introduced or removed amino acid residue comprising an attachment group for the non-polypeptide moiety. Conjugates of the present invention have a number of improved properties as compared to human interferon β , including reduced immunogenicity, increased functional *in vivo* half-life, increased serum half-life, and/or increased bioavailability. Consequently, the conjugate of the invention offers a number of advantages over the currently available interferon β compounds, including longer duration between injections, fewer side effects, and/or increased efficiency due to reduction in antibodies. Moreover, higher doses of active protein and thus a more effective therapeutic response may be

5 obtained by use of a conjugate of the invention. Furthermore, conjugates of the invention have demonstrated significantly reduced cross-reactivity with sera from patients treated with currently available interferon β products as defined hereinbelow.

In one aspect the invention relates to a conjugate exhibiting interferon β activity and comprising at least one first non-polypeptide moiety covalently attached to an interferon β polypeptide, the amino acid sequence of which differs from that of wild-type human interferon β in at least one introduced and at least one removed amino acid residue comprising an attachment group for said first non-polypeptide moiety.

In another aspect the invention relates to a conjugate exhibiting interferon β activity and comprising at least one first non-polypeptide moiety conjugated to at least one lysine residue of an interferon β polypeptide, the amino acid sequence of which differs from that of wild-type human interferon β in at least one introduced and/or at least one removed lysine residue.

In yet another aspect the invention relates to a conjugate exhibiting interferon β activity and comprising at least one first non-polypeptide moiety conjugated to at least one cysteine residue of an interferon β polypeptide, the amino acid sequence of which differs from at least one introduced cysteine residue into a position that in wild-type human interferon β is occupied by a surface exposed amino acid residue.

In yet another aspect the invention relates to a conjugate exhibiting interferon β activity and comprising at least one first non-polypeptide moiety having an acid group as an attachment group, which moiety is conjugated to at least one aspartic acid or glutamic acid residue of an interferon β polypeptide, the amino acid sequence of which differs from that of wild-type human interferon β in at least one introduced and/or at least one removed aspartic acid or glutamic acid residue.

In yet another aspect the invention relates to a conjugate exhibiting interferon β activity and comprising at least one polymer molecule and at least one sugar moiety covalently attached to an interferon β polypeptide, the amino acid sequence of which differs from that of wild-type human interferon β in

(a) at least one introduced and/or at least one removed amino acid residue comprising an attachment group for the polymer molecule, and

5 (b) at least one introduced and/or at least one removed amino acid residue comprising an attachment group for the sugar moiety,
provided that when the attachment group for the polymer molecule is a cysteine residue, and the sugar moiety is an N-linked sugar moiety, a cysteine residue is not inserted in such a manner that an N-glycosylation site is destroyed.

10 In yet another aspect the invention relates to a conjugate exhibiting interferon β activity and comprising an interferon β polypeptide, the amino acid sequence of which differs from that of wild-type human interferon β in at least one introduced glycosylation site, the conjugate further comprising at least one un-PEGylated sugar moiety attached to an introduced glycosylation site.

15 In yet another aspect the invention relates to a conjugate exhibiting interferon β activity and comprising an interferon β polypeptide, the amino acid sequence of which differs from that of wild-type human interferon β in that a glycosylation site has been introduced or removed by way of introduction or removal of amino acid residue(s) constituting a part of a glycosylation site in a position that in wildtype human interferon β is occupied by a surface exposed amino acid residue.

20 In a still further aspect the invention relates to a conjugate exhibiting interferon β activity and comprising a sugar moiety covalently attached to an interferon β polypeptide, the amino acid sequence of which differs from that of wild-type human interferon β in at least one removed glycosylation site.

25 In still further aspects the invention relates to means and methods for preparing a conjugate or interferon β polypeptide for use in the invention, including nucleotide sequences and expression vectors encoding the polypeptide as well as methods for preparing the polypeptide or the conjugate.

30 In final aspects the invention relates to a therapeutic composition comprising a conjugate of the invention, to a conjugate or composition of the invention for use in therapy, to the use of a conjugate or composition in therapy or for the manufacture of a medicament for treatment of diseases.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 illustrates the antiviral activity of a conjugate of the invention,

5 Figure 2 depicts the yield of interferon β production obtained according to
Example 8.

DETAILED DISCUSSION

DEFINITIONS

10 In the context of the present application and invention the following definitions
apply:

The term "conjugate" (or interchangeably "conjugated polypeptide") is intended
to indicate a heterogeneous (in the sense of composite or chimeric) molecule formed by the
covalent attachment of one or more polypeptide(s) to one or more non-polypeptide moieties. The
term covalent attachment means that the polypeptide and the non-polypeptide moiety are either
15 directly covalently joined to one another, or else are indirectly covalently joined to one another
through an intervening moiety or moieties, such as a bridge, spacer, or linkage moiety or
moieties using an attachment group present in the polypeptide. Preferably, the conjugate is
soluble at relevant concentrations and conditions, i.e. soluble in physiological fluids such as
blood. Examples of conjugated polypeptides of the invention include glycosylated and/or
20 PEGylated polypeptides. The term "non-conjugated polypeptide" may be used about the
polypeptide part of the conjugate.

The term "non-polypeptide moiety" is intended to indicate a molecule that is
capable of conjugating to an attachment group of the polypeptide of the invention. Preferred
examples of such molecule include polymer molecules, sugar moieties, lipophilic compounds, or
25 organic derivatizing agents. When used in the context of a conjugate of the invention it will be
understood that the non-polypeptide moiety is linked to the polypeptide part of the conjugate
through an attachment group of the polypeptide.

The term "polymer molecule" is defined as a molecule formed by covalent
linkage of two or more monomers, wherein none of the monomers is an amino acid residue,
30 except where the polymer is human albumin or another abundant plasma protein. The term
"polymer" may be used interchangeably with the term "polymer molecule". The term is intended
to cover carbohydrate molecules attached by *in vitro* glycosylation, i.e. a synthetic glycosylation
performed *in vitro* normally involving covalently linking a carbohydrate molecule to an
attachment group of the polypeptide, optionally using a cross-linking agent. Carbohydrate

5 molecules attached by *in vivo* glycosylation, such as N- or O-glycosylation (as further described below)) are referred to herein as “a sugar moiety”. Except where the number of non-polypeptide moieties, such as polymer molecule(s) or sugar moieties in the conjugate is expressly indicated every reference to “a non-polypeptide moiety” contained in a conjugate or otherwise used in the present invention shall be a reference to one or more non-polypeptide moieties, such as polymer
10 molecule(s) or sugar moieties, in the conjugate.

The term “attachment group” is intended to indicate an amino acid residue group of the polypeptide capable of coupling to the relevant non-polypeptide moiety. For instance, for polymer, in particular PEG, conjugation a frequently used attachment group is the ϵ -amino group of lysine or the N-terminal amino group. Other polymer attachment groups include a free
15 carboxylic acid group (e.g. that of the C-terminal amino acid residue or of an aspartic acid or glutamic acid residue), suitably activated carbonyl groups, oxidized carbohydrate moieties and mercapto groups.

For *in vivo* N-glycosylation, the term “attachment group” is used in an unconventional way to indicate the amino acid residues constituting an N-glycosylation site (with the sequence N-X'-S/T/C-X'', wherein X' is any amino acid residue except proline, X'' any amino acid residue that may or may not be identical to X' and preferably is different from proline, N is asparagine and S/T/C is either serine, threonine or cysteine, preferably serine or threonine, and most preferably threonine). Although the asparagine residue of the N-
25 glycosylation site is the one to which the sugar moiety is attached during glycosylation, such attachment cannot be achieved unless the other amino acid residues of the N-glycosylation site is present. Accordingly, when the non-polypeptide moiety is an N-linked sugar moiety, the term “amino acid residue comprising an attachment group for the non-polypeptide moiety” as used in connection with alterations of the amino acid sequence of the parent polypeptide is to be understood as amino acid residues constituting an N-glycosylation site is/are to be altered in such
30 a manner that either a functional N-glycosylation site is introduced into the amino acid sequence or removed from said sequence. For an “O-glycosylation site” the attachment group is the OH-group of a serine or threonine residue.

The term “one difference” or “differs from” as used in connection with specific mutations is intended to allow for additional differences being present apart from the specified
35 amino acid difference. For instance, in addition to the removal and/or introduction of amino acid

residues comprising an attachment group for the non-polypeptide moiety the interferon β polypeptide may comprise other substitutions that are not related to introduction and/or removal of such amino acid residues. The term “at least one” as used about a non-polypeptide moiety, an amino acid residue, a substitution, etc is intended to mean one or more. The terms “mutation” and “substitution” are used interchangeably herein.

In the present application, amino acid names and atom names (e.g. CA, CB, CD, CG, SG, NZ, N, O, C, etc) are used as defined by the Protein DataBank (PDB) (www.pdb.org) which are based on the IUPAC nomenclature (IUPAC Nomenclature and Symbolism for Amino Acids and Peptides (residue names, atom names e.t.c.), *Eur. J. Biochem.*, 138, 9-37 (1984) together with their corrections in *Eur. J. Biochem.*, 152, 1 (1985). CA is sometimes referred to as C α , CB as C β . The term “amino acid residue” is intended to indicate an amino acid residue contained in the group consisting of alanine (Ala or A), cysteine (Cys or C), aspartic acid (Asp or D), glutamic acid (Glu or E), phenylalanine (Phe or F), glycine (Gly or G), histidine (His or H), isoleucine (Ile or I), lysine (Lys or K), leucine (Leu or L), methionine (Met or M), asparagine (Asn or N), proline (Pro or P), glutamine (Gln or Q), arginine (Arg or R), serine (Ser or S), threonine (Thr or T), valine (Val or V), tryptophan (Trp or W), and tyrosine (Tyr or Y) residues. The terminology used for identifying amino acid positions/substitutions is illustrated as follows: C17 (indicates position #17 occupied by a cysteine residue in the amino acid sequence shown in SEQ ID NO 2). C17S (indicates that the cysteine residue of position 17 has been replaced with a serine). The numbering of amino acid residues made herein is made relative to the amino acid sequence shown in SEQ ID NO 2. “M1del” is used about a deletion of the methionine residue occupying position 1. Multiple substitutions are indicated with a “+”, e.g. R71N+D73T/S means an amino acid sequence which comprises a substitution of the arginine residue in position 71 with an asparagine and a substitution of the aspartic acid residue in position 73 with a threonine or serine residue, preferably a threonine residue. T/S as used about a given substitution herein means either a T or a S residue, preferably a T residue.

The term “nucleotide sequence” is intended to indicate a consecutive stretch of two or more nucleotide molecules. The nucleotide sequence may be of genomic, cDNA, RNA, semisynthetic, synthetic origin, or any combinations thereof.

The term “interferon β protein sequence family” is used in its conventional meaning, i.e. to indicate a group of polypeptides with sufficiently homologous amino acid

5 sequences to allow alignment of the sequences, e.g. using the CLUSTALW program. An
interferon β sequence family is available, e.g. from the PFAM families, version 4.0, or may be
prepared by use of a suitable computer program such as CLUSTALW version 1.74 using default
parameters (Thompson et al., 1994, CLUSTAL W: improving the sensitivity of progressive
multiple sequence alignment through sequence weighting, position-specific gap penalties and
10 weight matrix choice, Nucleic Acids Research, 22:4673-4680).

The term “polymerase chain reaction” or “PCR” generally refers to a method for
amplification of a desired nucleotide sequence in vitro, as described, for example, in US
4,683,195. In general, the PCR method involves repeated cycles of primer extension synthesis,
using oligonucleotide primers capable of hybridising preferentially to a template nucleic acid.

15 “Cell”, “host cell”, “cell line” and “cell culture” are used interchangeably herein
and all such terms should be understood to include progeny resulting from growth or culturing of
a cell. “Transformation” and “transfection” are used interchangeably to refer to the process of
introducing DNA into a cell.

20 “Operably linked” refers to the covalent joining of two or more nucleotide
sequences, by means of enzymatic ligation or otherwise, in a configuration relative to one
another such that the normal function of the sequences can be performed. For example, the
nucleotide sequence encoding a presequence or secretory leader is operably linked to a
nucleotide sequence for a polypeptide if it is expressed as a preprotein that participates in the
secretion of the polypeptide: a promoter or enhancer is operably linked to a coding sequence if it
25 affects the transcription of the sequence; a ribosome binding site is operably linked to a coding
sequence if it is positioned so as to facilitate translation. Generally, “operably linked” means that
the nucleotide sequences being linked are contiguous and, in the case of a secretory leader,
contiguous and in reading phase. Linking is accomplished by ligation at convenient restriction
sites. If such sites do not exist, then synthetic oligonucleotide adaptors or linkers are used, in
30 conjunction with standard recombinant DNA methods.

The term “introduce” is primarily intended to mean substitution of an existing
amino acid residue, but may also mean insertion of an additional amino acid residue. The term
“remove” is primarily intended to mean substitution of the amino acid residue to be removed by
another amino acid residue, but may also mean deletion (without substitution) of the amino acid
35 residue to be removed.

5 The term "immunogenicity" as used in connection with a given substance is intended to indicate the ability of the substance to induce a response from the immune system. The immune response may be a cell or antibody mediated response (see, e.g., Roitt: Essential Immunology (8th Edition, Blackwell) for further definition of immunogenicity). Immunogenicity may be determined by use of any suitable method known in the art, e.g. *in vivo* or *in vitro*, e.g. using the *in vitro* immunogenicity test outlined in the Materials and Methods section below. The term "reduced immunogenicity" is intended to indicate that the conjugate or polypeptide of the present invention gives rise to a measurably lower immune response than a reference molecule, such as wildtype human interferon β , e.g. Rebif or Avonex, or a variant of wild-type human interferon β such as Betaseron, as determined under comparable conditions. When reference is made herein to commercially available interferon β products (i.e. Betaseron, Avonex and Rebif), it should be understood to mean either the formulated product or the interferon β polypeptide part of the product (as appropriate). Normally, reduced antibody reactivity (e.g. reactivity towards antibodies present in serum from patients treated with commercial interferon β products) is an indication of reduced immunogenicity.

20 The term "functional *in vivo* half-life" is used in its normal meaning, i.e. the time at which 50% of a given functionality of the polypeptide or conjugate is retained (such as the time at which 50% of the biological activity of the polypeptide or conjugate is still present in the body/target organ, or the time at which the activity of the polypeptide or conjugate is 50% of the initial value). As an alternative to determining functional *in vivo* half-life, "serum half-life" may be determined, i.e. the time in which 50% of the polypeptide or conjugate molecules circulate in the plasma or bloodstream prior to being cleared. Determination of serum half-life is often more simple than determining functional *in vivo* half-life and the magnitude of serum half-life is usually a good indication of the magnitude of functional *in vivo* half-life. Alternative terms to serum half-life include "plasma half-life", "circulating half-life", "serum clearance", "plasma clearance" and "clearance half-life". The functionality to be retained is normally selected from antiviral, antiproliferative, immunomodulatory or receptor binding activity. Functional *in vivo* half-life and serum half-life may be determined by any suitable method known in the art as further discussed in the Materials and Methods section hereinafter.

35 The polypeptide or conjugate is normally cleared by the action of one or more of the reticuloendothelial systems (RES), kidney, spleen or liver, or by specific or unspecific

proteolysis. Clearance taking place by the kidneys may also be referred to as “renal clearance” and is e.g. accomplished by glomerular filtration, tubular excretion or tubular elimination. Normally, clearance depends on physical characteristics of the conjugate, including molecular weight, size (diameter) (relative to the cut-off for glomerular filtration), charge, symmetry, shape/rigidity, attached carbohydrate chains, and the presence of cellular receptors for the protein. A molecular weight of about 67 kDa is considered to be an important cut-off-value for renal clearance.

Reduced renal clearance may be established by any suitable assay, e.g. an established *in vivo* assay. Typically, the renal clearance is determined by administering a labelled (e.g. radiolabelled or fluorescence labelled) polypeptide conjugate to a patient and measuring the label activity in urine collected from the patient. Reduced renal clearance is determined relative to the corresponding non-conjugated polypeptide or the non-conjugated corresponding wild-type polypeptide or a commercial interferon β product under comparable conditions.

The term “increased” when used with respect to the functional *in vivo* half-life or serum half-life is used to indicate that the relevant half-life of the conjugate or polypeptide is statistically significantly increased relative to that of a reference molecule, such as an unconjugated wildtype human interferon β (e.g. Avonex™ or Rebif®) or an unconjugated variant human interferon β (e.g. Betaseron®) as determined under comparable conditions.

The term “reduced immunogenicity and/or increased functional *in vivo* half-life and/or increased serum half-life” is to be understood as covering any one, two or all of these properties. Preferably, a conjugate or polypeptide of the invention has at least two of these properties, i.e. reduced immunogenicity and increased functional *in vivo* half-life, reduced immunogenicity and increased serum half-life or increased functional *in vivo* half-life and increased serum half-life. Most preferably, the conjugate or polypeptide of the invention has all properties.

The term “exhibiting interferon β activity” is intended to indicate that the polypeptide or conjugate has one or more of the functions of native interferon β , in particular human wildtype interferon β with the amino acid sequence shown in SEQ ID NO 2 (which is the mature sequence) optionally expressed in a glycosylating host cell or any of the commercially available interferon β products. Such functions include capability to bind to an interferon receptor that is capable of binding interferon β and initiating intracellular signaling from the

receptor, in particular a type I interferon receptor constituted by the receptor subunits IFNAR-2 and IFNAR-1 (Domanski et al., The Journal of Biological Chemistry, Vol. 273, No. 6, pp3144-3147, 1998, Mogensen et al., Journal of Interferon and Cytokine Research, 19: 1069-1098, 1999), and antiviral, antiproliferative or immunomodulatory activity (which can be determined using assays known in the art (e.g. those cited in the following disclosure)). Interferon β activity may be assayed by methods known in the art as exemplified in the Materials and Methods section hereinafter.

The polypeptide or conjugate “exhibiting” or “having” interferon β activity is considered to have such activity, when it displays a measurable function, e.g. a measurable receptor binding and stimulating activity (e.g. as determined by the primary or secondary assay described in the Materials and Methods section). The polypeptide exhibiting interferon β activity may also be termed “interferon β molecule” or “interferon β polypeptide” herein. The term “interferon β polypeptide” is primarily used herein about modified polypeptides of the invention (having introduced or removed attachment groups for the relevant non-polypeptide moiety).

The term “parent interferon β ” is intended to indicate the starting molecule to be improved in accordance with the present invention. While the parent interferon β may be of any origin, such as vertebrate or mammalian origin (e.g. any of the origins defined in WO 00/23472), the parent interferon β is preferably wild-type human interferon β with SEQ ID NO 2 or a variant thereof. A “variant” is a polypeptide, which differs in one or more amino acid residues from a parent polypeptide, normally in 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acid residues. Examples of wild-type human interferon β include the polypeptide part of Avonex™ or Rebif®. An example of a parent interferon β variant is Betaseron®. Alternatively, the parent interferon β polypeptide may comprise an amino acid sequence, which is a hybrid molecule between interferon β and another homologous polypeptide, such as interferon α , optionally containing one or more additional substitutions introduced into the hybrid molecule. Such a hybrid molecule may contain an amino acid sequence, which differs in more than 10 amino acid residues from the amino acid sequence shown in SEQ ID NO 2. In order to be useful in the present invention the hybrid molecule exhibits interferon β activity (e.g. as determined in the secondary assay described in the Materials and Methods section herein).

5 The term "functional site" as used about a polypeptide or conjugate of the invention is intended to indicate one or more amino acid residues which is/are essential for or otherwise involved in the function or performance of interferon β , and thus "located at" the functional site. The functional site is e.g. a receptor binding site and may be determined by methods known in the art, preferably by analysis of a structure of the polypeptide complexed to a relevant receptor, such as the type I interferon receptor constituted by IFNAR-1 and IFNAR-2.

CONJUGATE OF THE INVENTION

As stated above, in a first aspect, the invention relates to a conjugate exhibiting interferon β activity and comprising at least one first non-polypeptide moiety covalently attached to an interferon β polypeptide, the amino acid sequence of which differs from that of wildtype human interferon β in at least one introduced and at least one removed amino acid residue comprising an attachment group for said first non-polypeptide moiety.

By removing and/or introducing amino acid residues comprising an attachment group for the non-polypeptide moiety it is possible to specifically adapt the polypeptide so as to make the molecule more susceptible to conjugation to the non-polypeptide moiety of choice, to optimize the conjugation pattern (e.g. to ensure an optimal distribution of non-polypeptide moieties on the surface of the interferon β molecule and thereby, e.g., effectively shield epitopes and other surface parts of the polypeptide without significantly impairing the function thereof). For instance, by introduction of attachment groups, the interferon β polypeptide is boosted or otherwise altered in the content of the specific amino acid residues to which the relevant non-polypeptide moiety binds, whereby a more efficient, specific and/or extensive conjugation is achieved. By removal of one or more attachment groups it is possible to avoid conjugation to the non-polypeptide moiety in parts of the polypeptide in which such conjugation is disadvantageous, e.g. to an amino acid residue located at or near a functional site of the polypeptide (since conjugation at such a site may result in inactivation or reduced interferon β activity of the resulting conjugate due to impaired receptor recognition). Further, it may be advantageous to remove an attachment group located closely to another attachment group in order to avoid heterogeneous conjugation to such groups.

It will be understood that the amino acid residue comprising an attachment group for a non-polypeptide moiety, either it be removed or introduced, is selected on the basis of the

Furthermore, in the interferon β polypeptide part of a conjugate of the invention attachment groups located at the receptor-binding site of interferon β has preferably been removed, preferably by substitution of the amino acid residue comprising such group.

A still further generally applicable approach for modifying an interferon β polypeptide is to shield, and thereby destroy or otherwise inactivate an epitope present in the parent interferon β , by conjugation to a non-polypeptide moiety. Epitopes of human interferon β may be identified by use of methods known in the art, also known as epitope mapping, see, e.g. Romagnoli et al., J. Biol Chem, 1999, 380(5):553-9, DeLisser HM, Methods Mol Biol, 1999, 96:11-20, Van de Water et al., Clin Immunol Immunopathol, 1997, 85(3):229-35, Saint-Remy JM, Toxicology, 1997, 119(1):77-81, and Lane DP and Stephen CW, Curr Opin Immunol, 1993, 5(2):268-71. One method is to establish a phage display library expressing random oligopeptides of e.g. 9 amino acid residues. IgG1 antibodies from specific antisera towards human interferon β are purified by immunoprecipitation and the reactive phages are identified by immunoblotting. By sequencing the DNA of the purified reactive phages, the sequence of the oligopeptide can be determined followed by localization of the sequence on the 3D-structure of the interferon β . Alternatively, epitopes can be identified according to the method described in US 5,041,376. The thereby identified region on the structure constitutes an epitope that then can be selected as a target region for introduction of an attachment group for the non-polypeptide moiety. Preferably, at least one epitope, such as two, three or four epitopes of human recombinant interferon β (optionally comprising the C17S mutation) are shielded by a non-polypeptide moiety according to the present invention. Accordingly, in one embodiment, the conjugate of the invention has at least one shielded epitope as compared to wild type human interferon β , optionally comprising the C17S mutation, including any commercially available interferon β . Preferably, the conjugate of the invention comprises a polypeptide that is modified so as to shield the epitope located in the vicinity of amino acid residue Q49 and/or F111. This may be done by introduction of an attachment group for a non-polypeptide moiety into a position located in the vicinity of (i.e. within 4 amino acid residues in the primary sequence or within about 10Å in the tertiary sequence) of Q49 and/or F111. The 10Å distance is measured between CB's (CA's in case of glycine). Such specific introductions are described in the following sections.

In case of removal of an attachment group, the relevant amino acid residue comprising such group and occupying a position as defined above is preferably substituted with a

different amino acid residue that does not comprise an attachment group for the non-polypeptide moiety in question.

In case of introduction of an attachment group, an amino acid residue comprising such group is introduced into the position, preferably by substitution of the amino acid residue occupying such position.

The exact number of attachment groups available for conjugation and present in the interferon β polypeptide is dependent on the effect desired to be achieved by conjugation. The effect to be obtained is, e.g., dependent on the nature and degree of conjugation (e.g. the identity of the non-polypeptide moiety, the number of non-polypeptide moieties desirable or possible to conjugate to the polypeptide, where they should be conjugated or where conjugation should be avoided, etc.). For instance, if reduced immunogenicity is desired, the number (and location of) attachment groups should be sufficient to shield most or all epitopes. This is normally obtained when a greater proportion of the interferon β polypeptide is shielded. Effective shielding of epitopes is normally achieved when the total number of attachment groups available for conjugation is in the range of 1-10 attachment groups, in particular in the range of 2-8, such as 3-7.

Functional *in vivo* half-life is e.g., dependent on the molecular weight of the conjugate and the number of attachment groups needed for providing increased half-life thus depends on the molecular weight of the non-polypeptide moiety in question. In one embodiment, the conjugate of the invention has a molecular weight of at least 67 kDa, in particular at least 70 kDa as measured by SDS-PAGE according to Laemmli, U.K., Nature Vol 227 (1970), p680-85. Interferon β has a molecular weight of about 20 kDa, and therefore additional about 50kDa is required to obtain the desired effect. This may be, e.g., be provided by 5 10kDa PEG molecules or as otherwise described herein.

In order to avoid too much disruption of the structure and function of the parent human interferon β molecule the total number of amino acid residues to be altered in accordance with the present invention (as compared to the amino acid sequence shown in SEQ ID NO 2) typically does not exceed 15. Preferably, the interferon β polypeptide comprises an amino acid sequence, which differs in 1-15 amino acid residues from the amino acid sequence shown in SEQ ID NO 2, such as in 1-8 or in 2-8 amino acid residues, e.g., in 1-5 or in 2-5 amino acid residues from the amino acid sequence shown in SEQ ID NO 2. Thus, normally the interferon β

polypeptide comprises an amino acid sequence that differs from the amino acid sequence shown in SEQ ID NO 2 in 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acid residues. Preferably, the above numbers represent either the total number of introduced or the total number of removed amino acid residues comprising an attachment group for the relevant non-polypeptide moiety, or the total number of introduced and removed amino acid residues comprising such group.

In the conjugate of the invention it is preferred that at least about 50% of all conjugatable attachment groups, such as at least about 80% and preferably all of such groups are occupied by the relevant non-polypeptide moiety. Accordingly, in a preferred embodiment the conjugate of the invention comprises, e.g., 1-10 non-polypeptide moieties, such as 2-8 or 3-6.

The conjugate of the invention has one or more of the following improved properties:

Reduced immunogenicity as compared to wild-type human interferon β (e.g. AvonexTM or Rebif[®]) or to Betaseron[®], e.g. a reduction of at least 25%, such as at least 50%, and more preferably at least 75%;

Increased functional *in vivo* half-life and/or increased serum half-life as compared to wild-type human interferon β (e.g. AvonexTM or Rebif[®]) or to Betaseron[®];

Reduced or no reaction with neutralizing antibodies from patients treated with wildtype human interferon β (e.g. Rebif[®] or AvonexTM) or with Betaseron[®], e.g. a reduction of neutralisation of at least 25%, such as of at least 50%, and preferably of at least 75%.

The magnitude of the antiviral activity of a conjugate of the invention may not be critical, and thus be reduced (e.g. by up to 75%) or increased (e.g. by at least 5%) or equal to that of wild-type human interferon β ((e.g. AvonexTM or Rebif[®]) or to Betaseron[®];

Furthermore, the degree of antiviral activity as compared to antiproliferative activity of a conjugate of the invention may vary, and thus be higher, lower or equal to that of wildtype human interferon β .

Conjugate of the invention, wherein the non-polypeptide moiety is a molecule that has lysine as an attachment group

In a preferred embodiment the first non-polypeptide moiety has lysine as an attachment group, and thus the interferon β polypeptide is one that comprises an amino acid sequence that differs from that of wildtype human interferon β in at least one introduced and/or

at least one removed lysine residue. While the non-polypeptide moiety may be any of those binding to a lysine residue, e.g. the ϵ -amino group thereof, such as a polymer molecule, a lipophilic group, an organic derivatizing agent or a carbohydrate moiety, it is preferably any of the polymer molecule mentioned in the section entitled "Conjugation to a polymer molecule", in particular a branched or linear PEG or polyalkylene oxide. Most preferably, the polymer molecule is PEG and the activated molecule to be used for conjugation is SS-PEG, NPC-PEG, aldehyd-PEG, mPEG-SPA, mPEG-SCM, mPEG-BTC from Shearwater Polymers, Inc, SC-PEG from Enzon, Inc., tresylated mPEG as described in US 5,880,255, or oxycarbonyl-oxy-N-dicarboxyimide-PEG (US 5,122,614). Normally, for conjugation to a lysine residue the non-polypeptide moiety has a molecular weight of about 5 or 10 kDa.

In one embodiment the amino acid sequence of the interferon β polypeptide differs from that of human wildtype interferon β in at least one removed lysine residue, such as 1-5 removed lysine residues, in particular 1-4 or 1-3 removed lysine residues. The lysine residue(s) to be removed, preferably by replacement, is selected from the group consisting of K19, K33, K45, K52, K99, K105, K108, K115, K123, K134, and K136. The lysine residue(s) may be replaced with any other amino acid residue, but is preferably replaced by an arginine or a glutamine residue in order to give rise to the least structural difference. In particular, the polypeptide part may be one, wherein K19, K45, K52 and/or K123, preferably K19, K45 and/or K123 has/have been replaced with another any other amino acid residue, preferably arginine or glutamine. For instance, the interferon β polypeptide part of a conjugate of the invention comprises a combination of amino acid substitutions selected from the following list:

K19R+K45R+K123R;
K19Q+K45R+K123R;
K19R+K45Q+K123R;
K19R+K45R+K123Q;
K19Q+K45Q+K123R;
K19R+K45Q+K123Q;
K19Q+K45R+K123Q;
K19Q+K45Q+K123Q;
K45R+K123R;
K45Q+K123R;
K45Q+K123Q;
K45R+K123Q;
K19R+K123R;
K19Q+K123R;
K19R+K123Q;

K19Q+K123Q;
K19R+K45R;
K19Q+K45R;
K19R+K45Q; or
K19Q+K45Q.

In addition or alternatively to the amino acid substitutions mentioned in the above list the polypeptide part may comprise at least one substitution selected from the group consisting of K33R, K33Q, K52R, K52Q, K99R, K99Q, K105R, K105Q, K108R, K108Q, K115R, K115Q, K134R, K134Q, K136R, and K136Q, e.g. at least one of the following substitutions:

K52R+K134R;
K99R+K136R;
K33R+K105R+K136R;
K52R+K108R+K134R;
K99R+K115R+K136R;
K19R+K33R+K45R+K123R;
K19R+K45R+K52R+K123R;
K19R+K33R+K45R+K52R+K123R; or
K19R+K45R+K52R+K99R+K123R.

In a further embodiment the the amino acid sequence of the interferon β polypeptide differs from that shown in SEQ ID NO 2 in that a lysine residue has been introduced by substitution of at least one amino acid residue occupying a position that in the parent interferon β molecule is occupied by a surface exposed amino acid residue, preferably an amino acid residue having at least 25%, such as at least 50% of its side chain exposed to the surface. Preferably, the amino acid residue to be substituted is selected from the group consisting of N4, F8, L9, R11, S12, F15, Q16, Q18, L20, W22, Q23, G26, R27, L28, E29, Y30, L32, R35, M36, N37, D39, P41, E42, E43, L47, Q48, Q49, T58, Q64, N65, F67, A68, R71, Q72, D73, S75, S76, G78, N80, E81, I83, E85, N86, A89, N90, Y92, H93, H97, T100, L102, E103, L106, E107, E109, D110, F111, R113, G114, L116, M117, L120, H121, R124, G127, R128, L130, H131, E137, Y138, H140, I145, R147, V148, E149, R152, Y155, F156, N158, R159, G162, Y163, R165 and N166 of SEQ ID NO 2.

More preferably, the amino acid sequence of the interferon β polypeptide differs from the amino acid sequence shown in SEQ ID NO 2 in that a lysine residue has been introduced, by substitution, of at least one amino acid residue occupying a position selected from the group consisting of N4, F8, L9, R11, S12, G26, R27, E29, R35, N37, D39, E42, L47, Q48, Q49, A68, R71, Q72, D73, S75, G78, N80, E85, N86, A89, Y92, H93, D110, F111, R113, L116,

- 5 Y+Z+K19R+K45Q+K123R;
Y+Z+K19R+K45R+K123Q;
Y+Z+K19Q+K45Q+K123R;
Y+Z+K19R+K45Q+K123Q;
Y+Z+K19Q+K45R+K123Q;
10 Y+Z+K19Q+K45Q+K123Q;
Y+Z+K45R+K123R;
Y+Z+K45Q+K123R;
Y+Z+K45Q+K123Q;
Y+Z+K45R+K123Q;
15 Y+Z+K19R+K123R;
Y+Z+K19Q+K123R;
Y+Z+K19R+K123Q;
Y+Z+K19Q+K123Q;
Y+Z+K19R+K45R;
20 Y+Z+K19Q+K45R;
Y+Z+K19R+K45Q; or
Y+Z+K19Q+K45Q, wherein Y is selected from the group of Q49K, F111K, N80K,
Q49K+F111K, Q49K+N80K, F111K+N80K and Q49K+F111K+N80K and Z is absent or
comprises at least one substitution selected from the group consisting of K33R, K33Q, K52R,
25 K52Q, K99R, K99Q, K105R, K105Q, K108R, K108Q, K115R, K115Q, K134R, K134Q,
K136R, and K136Q. Preferably, the interferon β polypeptide comprises the following
substitution Y+Z+K19R+K45Q+K123R, wherein Y and Z have the above meaning.

More specifically, according to this embodiment the interferon β polypeptide may
comprise one of the following substitutions:

- 30 K19R+K45R+F111K+K123R;
K19R+K45R+Q49K+F111K+K123R;
K19R+K45R+Q49K+K123R;
K19R+K45R+ F111K;
K19R+K45R+Q49K+F111K;
35 K19R+Q49K+K123R;
K19R+Q49K+F111K+K123R;
K45Q+F111K+K123Q;
K45R+Q49K+K123R; or
K45R+Q49K+F111K+K123R.

- 40 Especially for expression in a non-glycosylating host such as *E. coli* the interferon
 β polypeptide may contain the substitution N80K or C17S+N80K, optionally in combination
with one or more of K19R/Q; K45R/Q; K52R/Q or K123R/Q. The substitution N80K is of
particular interest, when the interferon β polypeptide is expressed in a non-glycosylating host

5 cell, since N80 constitutes part of an inherent glycosylation site of human interferon β and conjugation at such site may mimick natural glycosylation.

Furthermore, it is preferred that the conjugate according to this aspect comprises at least two first non-polypeptide moieties, such as 2-8 moieties. .

10 Conjugate of the invention wherein the non-polypeptide moiety binds to a cysteine residue

In a still further aspect, the invention relates a conjugate exhibiting interferon β activity and comprising at least one first non-polypeptide conjugated to at least one cysteine residue of an interferon β polypeptide, the amino acid sequence of which differs from that of wildtype human interferon β in that at least one cysteine residue has been introduced,
15 preferably by substitution, into a position that in the parent interferon β molecule is occupied by an amino acid residue that is exposed to the surface of the molecule, preferably one that has at least 25%, such as at least 50% of its side chain exposed to the surface. For instance, the amino acid residue is selected from the group consisting of F8, L9, R11, S12, F15, Q16, Q18, L20, W22, L28, L32, M36, P41, T58, Q64, N65, F67, I83, E85, N86, A89, N90, Y92, H93, H97,
20 T100, L102, E103, L106, M117, L120, H121, R124, G127, R128, L130, H131, H140, I145, R147, V148, E149, R152, Y155, and F156 of SEQ ID NO 2.

Additionally or alternatively, the substitution is preferably performed at a position occupied by a threonine or serine residue. For instance, such position is selected from the group consisting of S2, S12, S13, T58, S74, S75, S76, T77, T82, T100, T112, S118, S119, S139, T144,
25 and T161, more preferably S2, S12, S13, S74, S75, S76, T77, T82, T100, T112, S118, S119, S139, and T144 (side chain surface exposed), still more preferably S2, S12, S75, S76, T82, T100, S119 and S139 (at least 25% of its side chain exposed), and even more preferably S12, S75, T82 and T100 (at least 50% of its side chain exposed). Of the above threonine or serine substitutions, serine substitutions are preferred. Accordingly, in even more preferred
30 embodiments, the position is selected from the group consisting of S2, S12, S13, S74, S75, S76, S118, S119 and S139, more preferably S2, S12, S13, S74, S75, S76, S118, S119 and S139, even more preferably S2, S12, S75, S76, S119 and S139, and still more preferably S12 and S75.

In one embodiment, only one cysteine residue is introduced into the interferon β polypeptide in order to avoid formation of disulphide bridges between two or more introduced
35 cysteine residues. In this connection C17 present in wildtype human interferon β may be

5 removed, preferably by substitution, in particular by substitution with S or A. In another embodiment, two or more cysteine residues are introduced, such as 2-6 or 2-4 cysteine residues. Preferably, the interferon β polypeptide part of the conjugate according to this embodiment of the invention comprises the mutation L47C, Q48C, Q49C, D110C, F111C or R113C, in particular only one of these mutations, optionally in combination with the mutation C17S. Also,
10 the interferon β polypeptide may comprise the substitution C17S+N80C.

While the first non-polypeptide moiety according to this aspect of the invention may be any molecule which, when using the given conjugation method has cysteine as an attachment group (such as a carbohydrate moiety, a lipophilic group or an organic derivatizing agent), it is preferred that the non-polypeptide moiety is a polymer molecule. The polymer
15 molecule may be any of the molecules mentioned in the section entitled "Conjugation to a polymer molecule", but is preferably selected from the group consisting of linear or branched polyethylene glycol or polyalkylene oxide. Most preferably, the polymer molecule is VS-PEG. The conjugation between the polypeptide and the polymer may be achieved in any suitable manner, e.g. as described in the section entitled "Conjugation to a polymer molecule", e.g. in
20 using a one step method or in the stepwise manner referred to in said section. When the interferon β polypeptide comprises only one conjugatable cysteine residue, this is preferably conjugated to a first non-polypeptide moiety with a molecular weight of at least 20kDa, either directly conjugated or indirectly through a low molecular weight polymer (as disclosed in WO 99/55377). When the conjugate comprises two or more first non-polypeptide moieties, normally
25 each of these has a molecular weight of 5 or 10kDa.

Conjugate of the invention wherein the non-polypeptide moiety binds to an acid group

In a still further aspect the invention relates to a conjugate exhibiting interferon β activity and comprising at least one first non-polypeptide moiety having an acid group as the
30 attachment group, which moiety is conjugated to at least one aspartic acid residue or one glutamic acid residue of an interferon β polypeptide, the amino acid sequence of which differs from that of wildtype human interferon β in at least one introduced and/or at least one removed aspartic acid or glutamic acid residue, respectively. The relevant amino acid residue may be introduced in any position occupied by a surface exposed amino acid residue, preferably by an
35 amino acid residue having more than 25% of its side chain surface exposed. Preferably, at least

5 one amino acid residue occupying a position selected from the group consisting of N4, L5, L6, F8, L9, Q10, R11, S12, S13, F15, Q16, Q18, K19, L20, W22, Q23, L24, N25, G26, R27, Y30, M36, Q46, Q48, Q49, I66, F67, A68, I69, F70, R71, S75, T82, I83, L87, A89, N90, V91, Y92, H93, Q94, I95, N96, H97, K108, F111, L116, L120, K123, R124, Y126, G127, R128, L130, H131, Y132, K134, A135, H140, T144, R147, Y155, F156, N158, R159, G162, Y163 and R165
10 has been substituted with an aspartic acid residue or a glutamic acid residue.

More preferably, the position is selected from the group consisting of N4, L5, F8, L9, R11, S12, F15, Q16, Q18, K19, W22, Q23, G26, R27, Y30, M36, Q46, Q48, Q49, A68, R71, S75, T82, A89, N90, Y92, H93, N96, H97, K108, F111, L116, L120, K123, R124, G127, R128, L130, H131, K134, A135, H140, Y155, N158, R159, G162, Y163 and R165, such as from
15 the group consisting of N4, L5, F8, S12, F15, Q16, K19, W22, Q23, R27, Y30, M36, Q46, Q48, Q49, R71, S75, T82, A89, Y92, H93, K108, F111, L116, K123, R124, G127, H131, K134, A135, Y155 and R165, still more preferably from the group consisting of N4, L5, F8, S12, F15, Q16, K19, W22, Q23, R27, Y30, Q46, Q48, Q49, S75, T82, A89, Y92, H93, K108, F111, L116, R124, G127, H131, K134, Y155 and R165, such as from the group consisting of L5, F8, S12, F15, Q16, K19, W22, Q23, Q48, Q49, Y92, H93, R124, G127, H131 and Y155, even more
20 preferably from the group consisting of S12, Q16, K19, Q23, Q48, Q49, Y92, H93, R124, G127, H131 and Y155, such as from the group consisting of S12, Q16, K19, Q23, Q48, Y92, H93, R124, G127, H131 and Y155, in particular from the group consisting of S12, Q16, K19, Q23, Q48, H93 and H131, even more preferably from the group consisting of S12, Q16, K19, Q48, H93 and H131, and most preferably from the group consisting of Q16 and Q48.
25

Furthermore, in order to obtain a sufficient number of non-polypeptide moieties it is preferred that that least two aspartic acid residues or at least two glutamic acid residues be introduced, preferably in two positions selected from any of the above lists. Also, it is preferred that the conjugate according to this aspect comprises at least two first non-polypeptide moieties.

30 In case of removal of an amino acid residue, the amino acid sequence of the interferon β polypeptide differs from that of human wildtype interferon β in at least one removed aspartic acid or glutamic acid residue, such as 1-5 removed residues, in particular 1-4 or 1-3 removed aspartic acid or glutamic acid residues. The residue(s) to be removed, preferably by replacement, is selected from the group consisting of D34, D39, D54, D73, D110, E29, E42, E43, E53, E61, E81, E85, E103, E104, E107, E109, E137 and E149. The aspartic acid or
35

5 glutamic acid residue(s) may be replaced with any other amino acid residue, but is preferably replaced by an arginine or a glutamine residue. first non-polypeptide moiety can be any non-polypeptide moiety with such property, it is presently preferred that the non-polypeptide moiety is a polymer molecule or an organic derivatizing agent having an acid group as an attachment group, in particular a polymer molecule such as PEG, and the conjugate is prepared, e.g., as described by Sakane and Pardridge, Pharmaceutical Research, Vol. 14, No. 8, 1997, pp 1085-1091. Normally, for conjugation to an acid group the non-polypeptide moiety has a molecular weight of about 5 or 10 kDa.

Conjugate of the invention comprising a second non-polypeptide moiety
In addition to a first non-polypeptide moiety (as described in the preceding

15 sections), the conjugate of the invention may comprise a second non-polypeptide moiety of a different type as compared to the first non-polypeptide moiety. Preferably, in any of the above described conjugates wherein the first non-polypeptide moiety is, e.g., a polymer molecule such as PEG, a second non-polypeptide moiety is a sugar moiety, in particular an N-linked sugar moiety. While the second non-polypeptide moiety may be attached to a natural glycosylation site of human interferon β , e.g. the N-linked glycosylation site defined by N80, it is normally advantageous to introduce at least one additional glycosylation site in the interferon β polypeptide. Such site is e.g. any of those described in the immediately preceding section entitled "Conjugate of the invention wherein the non-polypeptide moiety is a sugar moiety". Furthermore, in case at least one additional glycosylation site is introduced this may be accompanied by removal of an existing glycosylation site as described below.

It will be understood that in order to obtain an optimal distribution of attached first and second non-polypeptide moieties, the interferon β polypeptide may be modified in the number and distribution of attachment groups for the first as well as the second non-polypeptide moiety so as to have e.g. at least one removed attachment group for the first non-polypeptide moiety and at least one introduced attachment group for the second non-polypeptide moiety or vice versa. For instance, the interferon β polypeptide comprises at least two (e.g. 2-5) removed attachment groups for the first non-polypeptide moiety and at least one (e.g. 1-5) introduced attachment groups for the second non-polypeptide moiety or vice versa. Of particular interest is a conjugate wherein the first non-polypeptide moiety is a polymer molecule such as PEG having

lysine as an attachment group, and the second non-polypeptide moiety is an N-linked sugar moiety.

More specifically, the conjugate of the invention may be one exhibiting interferon β activity and comprising at least one polymer molecule, preferably PEG, and at least one sugar moiety covalently attached to an interferon β polypeptide, the amino acid sequence of which differs from that of wild-type human interferon β in

a) at least one introduced and/or at least one removed amino acid residue comprising an attachment group for the polymer molecule; and
b) at least one introduced and/or at least one removed *in vivo* glycosylation site, in particular an N-glycosylation site,

provided that when the attachment group for the polymer molecule is a cysteine residue, and the sugar moiety is an N-linked sugar moiety, a cysteine residue is not inserted in such a manner that an N-glycosylation site is destroyed. WO 99/03887 suggests that a cysteine residue can be introduced into the natural N-glycosylation site of interferon β .

In a specific embodiment, the interferon β polypeptide comprises one of the following sets of mutations:

K19R+K45R+Q49N+Q51T+F111N+R113T+K123R;
K19R+K45R+Q49N+Q51T+F111N+R113T; or
K19R+K45R+Q49N+Q51T+ K123R.

Conjugate of the invention wherein the non-polypeptide moiety is a sugar moiety

When the conjugate of the invention comprises at least one sugar moiety attached to an *in vivo* glycosylation site, in particular an N-glycosylation site, this is either the natural N-glycosylation site of wild-type human interferon β at position N80, i.e. defined by amino acid residues N80, E81, T82 and I83, or a new *in vivo* glycosylation site introduced into the interferon β polypeptide. The *in vivo* glycosylation site may be an O-glycosylation site, but is preferably an N-glycosylation site.

More specifically, in one aspect the invention relates to a conjugate exhibiting interferon β activity and comprising an interferon β polypeptide, the amino acid sequence of which differs from that of wild-type human interferon β in at least one introduced glycosylation site, the conjugate further comprising at least one un-PEGylated sugar moiety attached to an introduced glycosylation site.

5 In another aspect the invention relates to a conjugate exhibiting interferon β activity and comprising an interferon β polypeptide, the amino acid sequence of which differs from that of wild-type human interferon β in that a glycosylation site has been introduced or removed, provided that if only a glycosylation site is removed (and thus that no glycosylation site is introduced) the interferon β polypeptide does not comprise one or more of the following
10 substitutions: N80C, E81C or T82C. The latter substitution is suggested in WO 99/03887.

For instance, an *in vivo* glycosylation site is introduced into a position of the parent interferon β molecule occupied by an amino acid residue exposed to the surface of the molecule, preferably with more than 25% of the side chain exposed to the solvent, in particular more than 50% exposed to the solvent (these positions are identified in the Methods section
15 herein). The N-glycosylation site is introduced in such a way that the N-residue of said site is located in said position. Analogously, an O-glycosylation site is introduced so that the S or T residue making up such site is located in said position. Furthermore, in order to ensure efficient glycosylation it is preferred that the *in vivo* glycosylation site, in particular the N residue of the N-glycosylation site or the S or T residue of the O-glycosylation site, is located within the first
20 141 amino acid residues of the interferon β polypeptide, more preferably within the first 116 amino acid residues. Still more preferably, the *in vivo* glycosylation site is introduced into a position wherein only one mutation is required to create the site (i.e. where any other amino acid residues required for creating a functional glycosylation site is already present in the molecule).

Substitutions that lead to introduction of an additional N-glycosylation site at
25 positions exposed at the surface of the interferon β molecule and occupied by amino acid residues having more than 25% of the side chain exposed to the surface include:
S2N+N4S/T, L6S/T, L5N+G7S/T, F8N+Q10S/T, L9N+R11S/T, R11N, R11N+S13T, S12N+N14S/T, F15N+C17S/T, Q16N+Q18S/T, Q18N+L20S/T, K19N+L21S/T, W22N+L24S/T, Q23N+H25S/T, G26N+L28S/T, R27N+E29S/T, L28S+Y30S/T, Y30N+L32S/T, L32N+D34S/T, K33N+R35S/T, R35N+N37S/T, M36N+F38S/T, D39S/T, D39N+P41S/T, E42N+I44S/T, Q43N+K45S/T, K45N+L47S/T, Q46N+Q48S/T, L47N+Q49T/S, Q48N+F50S/T, Q49N+Q51S/T, Q51N+E53S/T, K52N+D54S/T, L57N+I59S/T, Q64N+I66S/T, A68N+F70S/T, R71N+D73S/T, Q72N, Q72N+S74T, D73N, D73N+S75T, S75N+T77S, S75N, S76N+G78S/T, E81N+I83S/T, T82N+V84S/T, E85N+L87S/T, L88S/T, A89N+V91S/T, Y92S/T, Y92N+Q94S/T, H93N+I95S/T, L98S/T, H97N+K99S/T, K99N+V101S/T,
35

5 T100N+L102S/T, E103N+K105S/T, E104N+L106S/T, K105N+E107S/T, E107N+E109S/T,
K108N+D110S/T, E109N+F111S/T, D110N+T112S, D110N, F111N+R113S/T,
R113N+K115S/T, G114N+L116S/T, K115N+M117S/T, L116N, L116N+S118T,
S119N+H212S/T, L120N+L122S/T, H121N+K123S/T, K123N+Y125S/T, R124N+Y126S/T,
G127N+I129S/T, R128N+L130S/T, L130N+Y132S/T, H131N+L133S/T, K134N+K136S/T,
10 A135N+E137S/T, K136N+Y138S/T, E137N, Y138N+H140S/T, H140N+A142S/T,
V148N+I150S/T, R152N+F154S/T, Y155N+I157S/T, L160S/T, R159N+T161S, R159N,
G162N+L164S/T, and Y163N+R165S/T.

Substitutions that lead to introduction of an additional N-glycosylation site at
positions exposed at the surface of the interferon β molecule having more than 50% of the side
15 chain exposed to the surface include: L6S/T, L5N+G7S/T, F8N+Q10S/T, L9N+R11S/T,
S12N+N14S/T, F15N+C17S/T, Q16N+Q18S/T, K19N+L21S/T, W22N+L24S/T,
Q23N+H25S/T, G26N+L28S/T, R27N+E29S/T, Y30N+L32S/T, K33N+R35S/T,
R35N+N37S/T, M36N+F38S/T, D39S/T, D39N+P41S/T, E42N+I44S/T, Q46N+Q48S/T,
Q48N+F50S/T, Q49N+Q51S/T, Q51N+E53S/T, K52N+D54S/T, L57N+I59S/T, R71N+D73S/T,
20 D73N, D73N+S75T, S75N+T77S, S75N, S76N+G78S/T, E81N+I83S/T, T82N+V84S/T,
E85N+L87S/T, A89N+V91S/T, Y92S/T, Y92N+Q94S/T, H93N+I95S/T, T100N+L102S/T,
E103N+K105S/T, E104N+L106S/T, E107N+E109S/T, K108N+D110S/T, D110N+T112S,
D110N, F111N+R113S/T, R113N+K115S/T, L116N, L116N+S118T, K123N+Y125S/T,
R124N+Y126S/T, G127N+I129S/T, H131N+L133S/T, K134N+K136S/T, A135N+E137S/T,
25 E137N, V148N+I150S/T, and Y155N+I157S/T.

Among the substitutions mentioned in the above lists, those are preferred that
have the N residue introduced among the 141 N-terminal amino acid residues, in particular
among the 116 N-terminal amino acid residues.

Substitutions that lead to introduction of an N-glycosylation site by only one
30 amino acid substitution include: L6S/T, R11N, D39S/T, Q72N, D73N, S75N, L88S/T, Y92S/T,
L98S/T, D110N, L116N, E137N, R159N and L160S/T. Among these, a substitution is preferred
that is selected from the group consisting of L6S/T, R11N, D39S/T, Q72N, D73N, S75N,
L88S/T, Y92S/T, L98S/T, D110N and L116N, more preferably from the group consisting of
L6S/T, D39S/T, D73N, S75N, L88S/T, D110N, L116N and E137N; and most preferably

5 selected from the group consisting of L6S/T, D39S/T, D73N, S75N, L88S/T, D110N and L116N.

One especially preferred interferon β polypeptide according to this aspect include at least one of the following substitutions:

10 S2N+N4T/S, L9N+R11T/S, R11N, S12N+N14T/S, F15N+C17S/T, Q16N+Q18T/S, K19N+L21T/S, Q23N+H25T/S, G26N+L28T/S, R27N+E29T/S, L28N+Y30T/S, D39T/S, K45N+L47T/S, Q46N+Q48T/S, Q48N+F50T/S, Q49N+Q51T/S, Q51N+E53T/S, R71N+D73T/S, Q72N, D73N, S75N, S76N+G78T/S, L88T/S, Y92T/S, N93N+I95T/S, L98T/S, E103N+K105T/S, E104N+L106T/S, E107N+E109T/S, K108N+D110T/S, D110N, F111N+R113T/S, or L116N, more preferably at least one of the following substitutions:

15 S2N+N4T, L9N+R11T, 49N+Q51T or F111N+R113T or R71N+D73T, in particular 49N+Q51T or F111N+R113T or R71N+D73T. For instance, the interferon β polypeptide comprises one of the following sets of substitutions :

20 Q49N+Q51T+F111N+R113T ;
Q49N+Q51T+R71N+D73T+ F111N+ R113T ;
S2N+N4T+ F111N+R113T ;
S2N+N4T+Q49N+Q51T ;
S2N+N4T+Q49N+Q51T+F111N+R113T ;
S2N+N4T+L9N+R11T+Q49N+Q51T ;
S2N+N4T+L9N+R11T+F111N+R113T ;
25 S2N+N4T+L9N+R11T+Q49N+Q51T+F111N+R113T ;
L9N+R11T+Q49N+Q51T;
L9N+R11T+Q49N+Q51T+F111N+R113T ; or
L9N+R11T+F111N+R113T

30 It will be understood that in order to introduce a functional *in vivo* glycosylation site the amino acid residue inbetween the N-residue and the S/T residue is different from proline. Normally, the amino acid residue inbetween will be that occupying the relevant position in the amino acid sequence shown in SEQ ID NO 2. For instance, in the polypeptide comprising the substitutions Q49N+Q51S, position 50 is the position inbetween.

35 The interferon β polypeptide part of a conjugate of the invention may contain a single *in vivo* glycosylation site. However, in order to obtain efficient shielding of epitopes present on the surface of the parent polypeptide it is often desirable that the polypeptide comprises more than one *in vivo* glycosylation site, in particular 2-7 *in vivo* glycosylation sites, such as 2, 3, 4, 5, 6 or 7 *in vivo* glycosylation sites. Thus, the interferon β polypeptide may comprise one additional glycosylation site, or may comprise two, three, four, five, six, seven or

5 more introduced *in vivo* glycosylation sites, preferably introduced by one or more substitutions described in any of the above lists.

As indicated above, in addition to one or more introduced glycosylation sites, existing glycosylation sites may have been removed from the interferon β polypeptide. For instance, any of the above listed substitutions to introduce a glycosylation site may be combined
10 with a substitution to remove the natural N-glycosylation site of human wild-type interferon β . For instance, the interferon β polypeptide may comprise a substitution of N80, e.g. one of the substitutions N80K/C/D/E, when a first non-polypeptide polypeptide is one having one of K, C, D, E as an attachment group. For instance, the interferon β polypeptide may comprise at least one of the following substitutions: S2N+N4T/S, L9N+R11T/S, R11N, S12N+N14T/S,
15 F15N+C17S/T, Q16N+Q18T/S, K19N+L21T/S, Q23N+H25T/S, G26N+L28T/S, R27N+E29T/S, L28N+Y30T/S, D39T/S, K45N+L47T/S, Q46N+Q48T/S, Q48N+F50T/S, Q49N+Q51T/S, Q51N+E53T/S, R71N+D73T/S, Q72N, D73N, S75N, S76N+G78T/S, L88T/S, Y92T/S, N93N+I95T/S, L98T/S, E103N+K105T/S, E104N+L106T/S, E107N+E109T/S, K108N+D110T/S, D110N, F111N+R113T/S, or L116N in combination with N80K/C/D/E. More
20 specifically, the interferon β polypeptide may comprise the substitution: Q49N+Q51T or F111N+R113T or R71N+D73T, in particular Q49N+Q51T+F111N+R113T or Q49N+Q51T+R71N+D73T+ F111N+ R113T, in combination with N80K/C/D/E.

Any of the glycosylated variants disclosed in the present section having introduced and/or removed at least one glycosylation site, such as the variant comprising the
25 substitutions Q48N+F50T/S, Q48N+F50T/S+F111N+R113T/S, Q49N+Q51T/S, F111N+R113T/S, or Q49N+Q51T/S+F111N+R113T/S, may further be conjugated to a polymer molecule, such as PEG, or any other non-polypeptide moiety. For this purpose the conjugation may be achieved by use of attachment groups already present in the interferon β polypeptide or attachment groups may have been introduced and/or removed, in particular such that a total of 1-
30 6, in particular 3-4 or 1, 2, 3, 4, 5, or 6 attachment groups are available for conjugation. Preferably, in a conjugate of the invention wherein the interferon β polypeptide comprises two glycosylation sites, the number and molecular weight of the non-polypeptide moiety is chosen so as that the total molecular weight added by the non-polypeptide moiety is in the range of 20-40 kDa, in particular about 20 kDa or 30 kDa.

5 In particular, the glycosylated variant may be conjugated to a non-polypeptide moiety via a lysine attachment group, and one or more lysine residues of the parent polypeptide may have been removed, e.g. by any of the substitutions mentioned in the section entitled "Conjugate of the invention, wherein the non-polypeptide moiety is a molecule which has lysine as an attachment group", in particular the substitutions K19R+K45R+K123R. Alternatively or
10 additionally, a lysine residue may have been introduced, e.g. by any of the substitutions mentioned in said section, in particular the substitution R71K. Accordingly, one specific conjugate of the invention is one, which comprises a glycosylated interferon β polypeptide comprising the mutations Q49N + Q51T + F111N + R113T + K19R + K45R + K123R or Q49N + Q51T + F111N + R113T + K19R + K45R + K123R + R71K further conjugated to PEG. The
15 glycosylated polypeptide part of said conjugate is favourably produced in CHO cells and PEGylated subsequent to purification using e.g. SS-PEG, NPC-PEG, aldehyd-PEG, mPEG-SPA, mPEG-SCM, mPEG-BTC from Shearwater Polymers, Inc, SC-PEG from Enzon, Inc., tresylated mPEG as described in US 5,880,255, or oxycarbonyl-oxy-N-dicarboxyimide-PEG (US 5,122,614).

20 Alternatively, to PEGylation via a lysine group, the glycosylated conjugate according to this embodiment may be PEGylated via a cysteine group as described in the section entitled "Conjugate of the invention, wherein the non-polypeptide moiety is a molecule that has cysteine as an attachment group" (for this purpose the interferon β polypeptide may, e.g. comprising at least one of the mutations N80C, R71C and C17S), via an acid group as described
25 in the section entitled "Conjugation of the invention wherein the non-polypeptide moiety binds to an acid group", or via any other suitable group.

Other conjugates of the invention

In addition to the introduction and/or removal of amino acid residues comprising an attachment group for the non-polypeptide moiety of choice (as described in any of the
30 sections above entitled "Conjugate of the invention") the interferon β polypeptide part of the conjugate may contain further substitutions. A preferred example is a substitution of any of the residues, M1, C17, N80 or V101, e.g. one or more of the following substitutions: C17S; N80K/C/D/E; V101Y/W/F/,H; a deletion of M1; or M1K. The substitution M1K is of particular interest when the interferon β polypeptide is expressed with a tag, e.g. a His-14tag, where such
35 tag is to be removed by DAP (diaminopeptidase) subsequent to purification and/or conjugation.

Non-polypeptide moiety of the conjugate of the invention

As indicated further above the non-polypeptide moiety of the conjugate of the invention is preferably selected from the group consisting of a polymer molecule, a lipophilic compound, a sugar moiety (by way of *in vivo* glycosylation) and an organic derivatizing agent. All of these agents may confer desirable properties to the polypeptide part of the conjugate, in particular reduced immunogenicity and/or increased functional *in vivo* half-life and/or increased serum half-life. The polypeptide part of the conjugate may be conjugated to only one type of non-polypeptide moiety, but may also be conjugated to two or more different types of non-polypeptide moieties, e.g. to a polymer molecule and a sugar moiety, to a lipophilic group and a sugar moiety, to an organic derivating agent and a sugar moiety, to a lipophilic group and a polymer molecule, etc. The conjugation to two or more different non-polypeptide moieties may be done simultaneous or sequentially. The choice of non-polypeptide moiety/ies, e.g. depends on the effect desired to be achieved by the conjugation. For instance, sugar moieties have been found particularly useful for reducing immunogenicity, whereas polymer molecules such as PEG are of particular use for increasing functional *in vivo* half-life and/or serum half-life. Using a polymer molecule as a first non-polypeptide moiety and a sugar moiety as a second non-polypeptide moiety may result in reduced immunogenicity and increased functional *in vivo* or serum half-life.

Methods of preparing a conjugate of the invention

In the following sections "Conjugation to a lipophilic compound", "Conjugation to a polymer molecule", "Conjugation to a sugar moiety" and "Conjugation to an organic derivatizing agent" conjugation to specific types of non-polypeptide moieties is described.

Conjugation to a lipophilic compound

For conjugation to a lipophilic compound the following polypeptide groups may function as attachment groups: the N-terminal or C-terminal of the polypeptide, the hydroxy groups of the amino acid residues Ser, Thr or Tyr, the ϵ -amino group of Lys, the SH group of Cys or the carboxyl group of Asp and Glu. The polypeptide and the lipophilic compound may be conjugated to each other, either directly or by use of a linker. The lipophilic compound may be a natural compound such as a saturated or unsaturated fatty acid, a fatty acid diketone, a terpene, a prostaglandin, a vitamine, a carotenoide or steroide, or a synthetic compound such as a carbon acid, an alcohol, an amine and sulphonic acid with one or more alkyl-, aryl-, alkenyl- or other

multiple unsaturated compounds. The conjugation between the polypeptide and the lipophilic compound, optionally through a linker may be done according to methods known in the art, e.g. as described by Bodanszky in Peptide Synthesis, John Wiley, New York, 1976 and in WO 96/12505.

Conjugation to a polymer molecule

The polymer molecule to be coupled to the polypeptide may be any suitable polymer molecule, such as a natural or synthetic homo-polymer or heteropolymer, typically with a molecular weight in the range of 300-100,000 Da, such as 300-20,000 Da, more preferably in the range of 500-10,000 Da, even more preferably in the range of 500-5000 Da.

Examples of homo-polymers include a polyol (i.e. poly-OH), a polyamine (i.e. poly-NH₂) and a polycarboxylic acid (i.e. poly-COOH). A hetero-polymer is a polymer, which comprises one or more different coupling groups, such as, e.g., a hydroxyl group and an amine group.

Examples of suitable polymer molecules include polymer molecules selected from the group consisting of polyalkylene oxide (PAO), including polyalkylene glycol (PAG), such as polyethylene glycol (PEG) and polypropylene glycol (PPG), branched PEGs, poly-vinyl alcohol (PVA), poly-carboxylate, poly-(vinylpyrrolidone), polyethylene-co-maleic acid anhydride, polystyrene-co-malic acid anhydride, dextran including carboxymethyl-dextran, or any other biopolymer suitable for reducing immunogenicity and/or increasing functional *in vivo* half-life and/or serum half-life. Another example of a polymer molecule is human albumin or another abundant plasma protein. Generally, polyalkylene glycol-derived polymers are biocompatible, non-toxic, non-antigenic, non-immunogenic, have various water solubility properties, and are easily excreted from living organisms.

PEG is the preferred polymer molecule to be used, since it has only few reactive groups capable of cross-linking compared, e.g., to polysaccharides such as dextran, and the like. In particular, monofunctional PEG, e.g. monomethoxypolyethylene glycol (mPEG), is of interest since its coupling chemistry is relatively simple (only one reactive group is available for conjugating with attachment groups on the polypeptide). Consequently, the risk of cross-linking is eliminated, the resulting polypeptide conjugates are more homogeneous and the reaction of the polymer molecules with the polypeptide is easier to control.

5 To effect covalent attachment of the polymer molecule(s) to the polypeptide, the hydroxyl end groups of the polymer molecule must be provided in activated form, i.e. with reactive functional groups (examples of which include primary amino groups, hydrazide (HZ), thiol, succinate (SUC), succinimidyl succinate (SS), succinimidyl succinamide (SSA), succinimidyl propionate (SPA), succinimidy carboxymethylate (SCM), benzotriazole carbonate (BTC), N-hydroxysuccinimide (NHS), aldehyde, nitrophenylcarbonate (NPC), and tresylate (TRES)). Suitably activated polymer molecules are commercially available, e.g. from Shearwater Polymers, Inc., Huntsville, AL, USA. Alternatively, the polymer molecules can be activated by conventional methods known in the art, e.g. as disclosed in WO 90/13540. Specific examples of activated linear or branched polymer molecules for use in the present invention are described in the Shearwater Polymers, Inc. 1997 and 2000 Catalogs (Functionalized Biocompatible Polymers for Research and pharmaceuticals, Polyethylene Glycol and Derivatives, incorporated herein by reference). Specific examples of activated PEG polymers include the following linear PEGs: NHS-PEG (e.g. SPA-PEG, SSPA-PEG, SBA-PEG, SS-PEG, SSA-PEG, SC-PEG, SG-PEG, and SCM-PEG), and NOR-PEG), BTC-PEG, EPOX-PEG, NCO-PEG, NPC-PEG, CDI-PEG, ALD-PEG, TRES-PEG, VS-PEG, IODO-PEG, and MAL-PEG, and branched PEGs such as PEG2-NHS and those disclosed in US 5,932,462 and US 5,643,575, both of which references are incorporated herein by reference. Furthermore, the following publications, incorporated herein by reference, disclose useful polymer molecules and/or PEGylation chemistries: US 5,824,778, US 5,476,653, WO 97/32607, EP 229,108, EP 402,378, US 4,902,502, US 5,281,698, US 5,122,614, US 5,219,564, WO 92/16555, WO 94/04193, WO 94/14758, WO 94/17039, WO 94/18247, WO 94/28024, WO 95/00162, WO 95/11924, WO95/13090, WO 95/33490, WO 96/00080, WO 97/18832, WO 98/41562, WO 98/48837, WO 99/32134, WO 99/32139, WO 99/32140, WO 96/40791, WO 98/32466, WO 95/06058, EP 439 508, WO 97/03106, WO 96/21469, WO 95/13312, EP 921 131, US 5,736,625, WO 98/05363, EP 809 996, US 5,629,384, WO 96/41813, WO 96/07670, US 5,473,034, US 5,516,673, EP 605 963, US 5,382,657, EP 510 356, EP 400 472, EP 183 503 and EP 154 316.

The conjugation of the polypeptide and the activated polymer molecules is conducted by use of any conventional method, e.g. as described in the following references (which also describe suitable methods for activation of polymer molecules): Harris and Zalipsky, eds., Poly(ethylene glycol) Chemistry and Biological Applications, AZC, Washington; R.F.

5 Taylor, (1991), "Protein immobilisation. Fundamental and applications", Marcel Dekker, N.Y.;
S.S. Wong, (1992), "Chemistry of Protein Conjugation and Crosslinking", CRC Press, Boca
Raton; G.T. Hermanson et al., (1993), "Immobilized Affinity Ligand Techniques", Academic
Press, N.Y.). The skilled person will be aware that the activation method and/or conjugation
chemistry to be used depends on the attachment group(s) of the interferon β polypeptide as well
10 as the functional groups of the polymer (e.g. being amino, hydroxyl, carboxyl, aldehyde or sulfy-
dryl). The PEGylation may be directed towards conjugation to all available attachment groups on
the polypeptide (i.e. such attachment groups that are exposed at the surface of the polypeptide) or
may be directed towards specific attachment groups, e.g. the N-terminal amino group (US
5,985,265). Furthermore, the conjugation may be achieved in one step or in a stepwise manner
15 (e.g. as described in WO 99/55377).

It will be understood that the PEGylation is designed so as to produce the optimal
molecule with respect to the number of PEG molecules attached, the size and form (e.g. whether
they are linear or branched) of such molecules, and where in the polypeptide such molecules are
attached. For instance, the molecular weight of the polymer to be used may be chosen on the basis
20 of the desired effect to be achieved. For instance, if the primary purpose of the conjugation is to
achieve a conjugate having a high molecular weight (e.g. to reduce renal clearance) it is usually
desirable to conjugate as few high Mw polymer molecules as possible to obtain the desired
molecular weight. When a high degree of epitope shielding is desirable this may be obtained by use
of a sufficiently high number of low molecular weight polymer (e.g. with a molecular weight of
25 about 5,000 Da) to effectively shield all or most epitopes of the polypeptide. For instance, 2-8, such
as 3-6 such polymers may be used.

In connection with conjugation to only a single attachment group on the protein
(as described in US 5,985,265), it may be advantageous that the polymer molecule, which may
be linear or branched, has a high molecular weight, e.g. about 20 kDa.

30 Normally, the polymer conjugation is performed under conditions aiming at
reacting all available polymer attachment groups with polymer molecules. Typically, the molar
ratio of activated polymer molecules to polypeptide is 1000-1, in particular 200-1, preferably
100-1, such as 10-1 or 5-1 in order to obtain optimal reaction. However, also equimolar ratios
may be used.

5 It is also contemplated according to the invention to couple the polymer molecules to the polypeptide through a linker. Suitable linkers are well known to the skilled person. A preferred example is cyanuric chloride (Abuchowski et al., (1977), J. Biol. Chem., 252, 3578-3581; US 4,179,337; Shafer et al., (1986), J. Polym. Sci. Polym. Chem. Ed., 24, 375-378.

10 Subsequent to the conjugation residual activated polymer molecules are blocked according to methods known in the art, e.g. by addition of primary amine to the reaction mixture, and the resulting inactivated polymer molecules removed by a suitable method.

Covalent *in vitro* coupling of a carbohydrate moiety to amino acid residues of interferon β may be used to modify or increase the number or profile of carbohydrate substituents. Depending on the coupling mode used, the carbohydrate(s) may be attached to a) arginine and histidine (Lundblad and Noyes, Chemical Reagents for Protein Modification, CRC Press Inc. Boca Raton, FI), b) free carboxyl groups (e.g. of the C-terminal amino acid residue, asparagine or glutamine), c) free sulfhydryl groups such as that of cysteine, d) free hydroxyl groups such as those of serine, threonine, tyrosine or hydroxyproline, e) aromatic residues such as those of phenylalanine or tryptophan or f) the amide group of glutamine. These amino acid residues constitute examples of attachment groups for a carbohydrate moiety, which may be introduced and/or removed in the interferon β polypeptide. Suitable methods of *in vitro* coupling are described in WO 87/05330 and in Aplin et al., CRC Crit Rev. Biochem., pp. 259-306, 1981. The *in vitro* coupling of carbohydrate moieties or PEG to protein- and peptide-bound Gln-residues can also be carried out by transglutaminases (TGases), e.g. as described by Sato et al., 1996 Biochemistry 35, 13072-13080 or in EP 725145

Coupling to a sugar moiety

In order to achieve *in vivo* glycosylation of an interferon β polypeptide that has been modified by introduction of one or more glycosylation sites (see the section "Conjugates of the invention wherein the non-polypeptide moiety is a sugar moiety"), the nucleotide sequence encoding the polypeptide part of the conjugate must be inserted in a glycosylating, eucaryotic expression host. The expression host cell may be selected from fungal (filamentous fungal or yeast), insect, mammalian animal cells, from transgenic plant cells or from transgenic animals. Furthermore, the glycosylation may be achieved in the human body when using a nucleotide sequence encoding the polypeptide part of a conjugate of the invention or a polypeptide of the

invention in gene therapy. In one embodiment the host cell is a mammalian cell, such as an CHO cell, BHK or HEK cell, e.g. HEK293, or an insect cell, such as an SF9 cell, or a yeast cell, e.g. *Saccharomyces cerevisiae*, *Pichia pastoris* or any other suitable glycosylating host, e.g. as described further below. Optionally, sugar moieties attached to the interferon β polypeptide by *in vivo* glycosylation are further modified by use of glycosyltransferases, e.g. using the glycoAdvance™ technology marketed by Neose, Horsham, PA, USA. Thereby, it is possible to, e.g., increase the sialylation of the glycosylated interferon β polypeptide following expression and *in vivo* glycosylation by CHO cells.

Coupling to an organic derivatizing agent

Covalent modification of the interferon β polypeptide may be performed by reacting (an) attachment group(s) of the polypeptide with an organic derivatizing agent. Suitable derivatizing agents and methods are well known in the art. For example, cysteinyl residues most commonly are reacted with α -haloacetates (and corresponding amines), such as chloroacetic acid or chloroacetamide, to give carboxymethyl or carboxyamidomethyl derivatives. Cysteinyl residues also are derivatized by reaction with bromotrifluoroacetone, α -bromo- β -(4-imidozoyl)propionic acid, chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl 2-pyridyl disulfide, p-chloromercuribenzoate, 2-chloromercuri-4-nitrophenol, or chloro-7-nitrobenzo-2-oxa-1,3-diazole. Histidyl residues are derivatized by reaction with diethylpyrocarbonate at pH 5.5-7.0 because this agent is relatively specific for the histidyl side chain. Para-bromophenacyl bromide also is useful; the reaction is preferably performed in 0.1 M sodium cacodylate at pH 6.0. Lysinyl and amino terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysinyl residues. Other suitable reagents for derivatizing α -amino-containing residues include imidoesters such as methyl picolinimate; pyridoxal phosphate; pyridoxal; chloroborohydride; trinitrobenzenesulfonic acid; O-methylisourea; 2,4-pentanedione; and transaminase-catalyzed reaction with glyoxylate. Arginyl residues are modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3-butanedione, 1,2-cyclohexanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high pKa of the guanidine functional group. Furthermore, these reagents may react with the groups of lysine as well as the arginine guanidino group. Carboxyl side groups (aspartyl or glutamyl or C-terminal amino acid residue) are

5 selectively modified by reaction with carbodiimides ($R-N=C=N-R'$), where R and R' are different alkyl groups, such as 1-cyclohexyl-3-(2-morpholinyl-4-ethyl) carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl) carbodiimide. Furthermore, aspartyl and glutamyl residues are converted to asparaginyll and glutaminyll residues by reaction with ammonium ions.

Blocking of functional site

10 It has been reported that excessive polymer conjugation can lead to a loss of activity of the interferon β polypeptide to which the polymer is conjugated. This problem can be eliminated, e.g., by removal of attachment groups located at the functional site or by blocking the functional site prior to conjugation. These latter strategies constitute further embodiments of the invention (the first strategy being exemplified further above, e.g. by removal of lysine residues
15 which may be located close to a functional site). More specifically, according to the second strategy the conjugation between the interferon β polypeptide and the non-polypeptide moiety is conducted under conditions where the functional site of the polypeptide is blocked by a helper molecule capable of binding to the functional site of the polypeptide. Preferably, the helper molecule is one, which specifically recognizes a functional site of the polypeptide, such as a
20 receptor, in particular the type I interferon receptor. Alternatively, the helper molecule may be an antibody, in particular a monoclonal antibody recognizing the interferon β polypeptide. In particular, the helper molecule may be a neutralizing monoclonal antibody.

The polypeptide is allowed to interact with the helper molecule before effecting conjugation. This ensures that the functional site of the polypeptide is shielded or protected and
25 consequently unavailable for derivatization by the non-polypeptide moiety such, as a polymer. Following its elution from the helper molecule, the conjugate between the non-polypeptide moiety and the polypeptide can be recovered with at least a partially preserved functional site.

The subsequent conjugation of the polypeptide having a blocked functional site to a polymer, a lipophilic compound, an organic derivatizing agent or any other compound is
30 conducted in the normal way, e.g. as described in the sections above entitled "Conjugation to".

Irrespective of the nature of the helper molecule to be used to shield the functional site of the polypeptide from conjugation, it is desirable that the helper molecule is free from or comprises only a few attachment groups for the non-polypeptide moiety of choice in part(s) of
35 the molecule, where the conjugation to such groups will hamper the desorption of the conjugated

5 polypeptide from the helper molecule. Hereby, selective conjugation to attachment groups present in non-shielded parts of the polypeptide can be obtained and it is possible to reuse the helper molecule for repeated cycles of conjugation. For instance, if the non-polypeptide moiety is a polymer molecule such as PEG, which has the epsilon amino group of a lysine or N-terminal amino acid residue as an attachment group, it is desirable that the helper molecule is substantially
10 free from conjugatable epsilon amino groups, preferably free from any epsilon amino groups. Accordingly, in a preferred embodiment the helper molecule is a protein or peptide capable of binding to the functional site of the polypeptide, which protein or peptide is free from any conjugatable attachment groups for the non-polypeptide moiety of choice.

In a further embodiment, the helper molecule is first covalently linked to a solid
15 phase such as column packing materials, for instance Sephadex or agarose beads, or a surface, e.g. reaction vessel. Subsequently, the polypeptide is loaded onto the column material carrying the helper molecule and conjugation carried out according to methods known in the art, e.g. as described in the sections above entitled "Conjugation to". This procedure allows the polypeptide conjugate to be separated from the helper molecule by elution. The polypeptide
20 conjugate is eluted by conventional techniques under physico-chemical conditions that do not lead to a substantive degradation of the polypeptide conjugate. The fluid phase containing the polypeptide conjugate is separated from the solid phase to which the helper molecule remains covalently linked. The separation can be achieved in other ways: For instance, the helper molecule may be derivatised with a second molecule (e.g. biotin) that can be recognized by a
25 specific binder (e.g. streptavidin). The specific binder may be linked to a solid phase thereby allowing the separation of the polypeptide conjugate from the helper molecule-second molecule complex through passage over a second helper-solid phase column which will retain, upon subsequent elution, the helper molecule-second molecule complex, but not the polypeptide conjugate. The polypeptide conjugate may be released from the helper molecule in any
30 appropriate fashion. De-protection may be achieved by providing conditions in which the helper molecule dissociates from the functional site of the interferon β to which it is bound. For instance, a complex between an antibody to which a polymer is conjugated and an anti-idiotypic antibody can be dissociated by adjusting the pH to an acid or alkaline pH.

Conjugation of a tagged interferon β polypeptide

In an alternative embodiment the interferon β polypeptide is expressed, as a fusion protein, with a tag, i.e. an amino acid sequence or peptide stretch made up of typically 1-30, such as 1-20 or 1-15 or 1-10 amino acid residues. Besides allowing for fast and easy purification, the tag is a convenient tool for achieving conjugation between the tagged polypeptide and the non-polypeptide moiety. In particular, the tag may be used for achieving conjugation in microtiter plates or other carriers, such as paramagnetic beads, to which the tagged polypeptide can be immobilised via the tag. The conjugation to the tagged polypeptide in, e.g., microtiter plates has the advantage that the tagged polypeptide can be immobilised in the microtiter plates directly from the culture broth (in principle without any purification) and subjected to conjugation. Thereby, the total number of process steps (from expression to conjugation) can be reduced. Furthermore, the tag may function as a spacer molecule ensuring an improved accessibility to the immobilised polypeptide to be conjugated. The conjugation using a tagged polypeptide may be to any of the non-polypeptide moieties disclosed herein, e.g. to a polymer molecule such as PEG.

The identity of the specific tag to be used is not critical as long as the tag is capable of being expressed with the polypeptide and is capable of being immobilised on a suitable surface or carrier material. A number of suitable tags are commercially available, e.g. from Unizyme Laboratories, Denmark. For instance, the tag may be any of the following sequences:

His-His-His-His-His-His
Met-Lys-His-His-His-His-His
Met-Lys-His-His-Ala-His-His-Gln-His-His
Met-Lys-His-Gln-His-Gln-His-Gln-His-Gln-His-Gln
(vectors useful for providing such tags are available from Unizyme Laboratories, Denmark)

or any of the following:

EQKLI SEEDL (a C-terminal tag described in Mol. Cell. Biol. 5:3610-16, 1985)
DYKDDDDK (a C- or N-terminal tag)
YPYDVPDYA

Antibodies against the above tags are commercially available, e.g. from ADI,

Aves Lab and Research Diagnostics.

A convenient method for using a tagged polypeptide for PEGylation is given in the Materials and Methods section below.

5 The subsequent cleavage of the tag from the polypeptide may be achieved by use of commercially available enzymes.

Polypeptides of the Invention

In further aspects the invention relates to generally novel interferon β polypeptides described herein that, as compared to human wildtype interferon β has at least one introduced and/or at least one removed attachment group for a non-polypeptide moiety. The novel polypeptides are important intermediate compounds for the preparation of a conjugate of the invention. In addition, the polypeptides themselves may have interesting properties.

Examples of such polypeptides include those that comprises an amino acid sequence which differs from that of wild-type human interferon β in that at least one amino acid residue selected from the group consisting of N4, F8, L9, Q10, R11, S13, L24, N25, G26, L28, E29, N37, F38, Q48, Q49, Q64, N65, I66, F67, A68, I69, F70, R71, Q72, D73, S74, S75, S76, T77, G78, W79, N80, E81, T82, I83, V84, L87, L88, A89, N90, V91, Y92, H93, Q94, D110, F111, T112, R113, R128, H140, T144, I145, R147, V148, L151, R152, F154, Y155, N158 and N166 is replaced with a different amino acid residue selected from the group consisting of K, R, D, E, C and N. The amino acid residues specified above are located in positions, which are exposed at the surface of human interferon β molecule as demonstrated by the solved 3D structure of human interferon β . By replacing one or more of these residue with either of K, R, D, E, C and N attachment group(s) for a non-polypeptide moiety, in particular a polymer attachment group or an amino acid residue susceptible to modification by a carbohydrate moiety, is/are introduced into human interferon β . The resulting modified human interferon β molecule is a suitable starting compound for the preparation of an interferon β conjugate having improved properties as compared to the unmodified human interferon β molecule.

In a further aspect the invention relates to an interferon β polypeptide comprising an amino acid sequence which differs from that of wild-type human interferon β in that at least one amino acid residue selected from the group consisting of N4, F8, L9, Q10, R11, S12, S13, L24, N25, G26, L28, E29, N37, F38, D39, Q48, Q49, Q64, N65, I66, F67, A68, I69, F70, R71, Q72, D73, S74, S75, S76, T77, G78, W79, N80, E81, T82, I83, V84, E85, L87, L88, A89, N90, V91, Y92, H93, Q94, D110, F111, T112, R113, R128, H140, T144, I145, R147, V148, L151, R152, F154, Y155, N158, G162, and N166 is replaced with a lysine residue, provided that the polypeptide is different from the one having the amino acid sequence of wild-type human

5 interferon β with the following substitutions: D54N+E85K+V91I+V101M and different from one which is a hybrid molecule between interferon β and interferon α which as a consequence of being a hybrid has a lysine in position 39. The first of the disclaimed polypeptides is disclosed by Stewart et al, DNA Vol 6 no2 1987 p119-128 and was found to be inactive, the second is disclosed in US 4,769,233 and was constructed with the purpose of improving the biological
10 activity of interferon β . None of the disclaimed polypeptides were made for or described as being suitable intermediates for the preparation of interferon β conjugates with reduced immunogenicity and/or prolonged functional *in vivo* half-life and/or serum half-life.

A still further example includes an interferon β polypeptide comprising an amino acid sequence which differs from that of SEQ ID NO 2 in one or more substitutions selected
15 from the group consisting of N4K, F15K, Q16K, R27K, R35K, D39K, Q49K, E85K, A89K, E103K, E109K, R124K, E137K and R159K, provided that when the substitution is R27K the polypeptide is different from the one having the amino acid sequence of wild-type human interferon β with the following substitutions: R27K+E43K. The disclaimed polypeptide is disclosed by Stewart et al, DNA Vol 6 no2 1987 p119-128 and was found to have a low activity.
20 The polypeptide was made in the course of a study of function-structure relationship and was not mentioned as a possible intermediate product for the preparation of improved interferon β conjugate molecules. For instance, the interferon β polypeptide comprises an amino acid sequence, which differs from that of SEQ ID NO 2 in that it comprises the substitution R27K in combination with at least one additional substitution that is different from E43K, or the
25 substitution R35K in combination with at least one additional substitution provided that the polypeptide has an amino acid sequence which is different from the amino acid sequence of wild-type human interferon β modified with the following substitutions:
G7E+S12N+C17Y+R35K. The disclaimed polypeptide is disclosed by Stewart et al, DNA Vol 6
no2 1987 p119-128 as having a retained antiproliferative activity on Daudi cells relative to their
30 antiviral activity, but reduced overall activity as compared to wild type interferon β . The disclaimed polypeptide was not prepared with the purpose of reducing the immunogenicity and/or increasing the functional *in vivo* half-life and/or serum half-life, but was made in the course of a study of the structural functional relationship of interferon β .

5 R27K+K123R+R159K;
 R27K+K45R+K123R+R159K;
 R27K+Q49K+E85K+A89K+K123R;
 R27K+K45R+Q49K+E85K+A89K+K123R;
 R27K+D39K+Q49K+E85K+A89K+K123R;
 10 R27K+D39K+K45R+Q49K+E85K+A89K+K123R;
 N4K+R27K+D39K+Q49K+E85K+A89K+K123R; and
 N4K+R27K+D39K+K45R+Q49K+E85K+A89K+K123R.

It will be understood that any of the polypeptides of the invention disclosed herein may be used to prepare a conjugate of the invention, i.e. be covalently coupled to any of the non-polypeptide moieties disclosed herein. In particular, when a polypeptide of the invention is expressed in a glycosylating microorganism the polypeptide may be provided in glycosylated form.

Methods of preparing an interferon β polypeptide for use in the invention

The polypeptide of the present invention or the polypeptide part of a conjugate of the invention, optionally in glycosylated form, may be produced by any suitable method known in the art. Such methods include constructing a nucleotide sequence encoding the polypeptide and expressing the sequence in a suitable transformed or transfected host. However, polypeptides of the invention may be produced, albeit less efficiently, by chemical synthesis or a combination of chemical synthesis or a combination of chemical synthesis and recombinant DNA technology.

The nucleotide sequence of the invention encoding an interferon β polypeptide may be constructed by isolating or synthesizing a nucleotide sequence encoding the parent interferon β , e.g. with the amino acid sequence shown in SE Q ID NO 2, and then changing the nucleotide sequence so as to effect introduction (i.e. insertion or substitution) or deletion (i.e. removal or substitution) of the relevant amino acid residue(s).

The nucleotide sequence is conveniently modified by site-directed mutagenesis in accordance with well-known methods, see, e.g., Mark et al., "Site-specific Mutagenesis of the Human Fibroblast Interferon Gene", Proc. Natl. Acad. Sci. USA, 81, pp. 5662-66 (1984); and US 4,588,585.

Alternatively, the nucleotide sequence is prepared by chemical synthesis, e.g. by using an oligonucleotide synthesizer, wherein oligonucleotides are designed based on the amino acid sequence of the desired polypeptide, and preferably selecting those codons that are favored in the host cell in which the recombinant polypeptide will be produced. For example, several small oligonucleotides coding for portions of the desired polypeptide may be synthesized and

5 assembled by PCR, ligation or ligation chain reaction (LCR). The individual oligonucleotides typically contain 5' or 3' overhangs for complementary assembly.

Once assembled (by synthesis, site-directed mutagenesis or another method), the nucleotide sequence encoding the interferon β polypeptide is inserted into a recombinant vector and operably linked to control sequences necessary for expression of the interferon β in the
10 desired transformed host cell.

It should of course be understood that not all vectors and expression control sequences function equally well to express the nucleotide sequence encoding a polypeptide variant described herein. Neither will all hosts function equally well with the same expression system. However, one of skill in the art may make a selection among these vectors, expression
15 control sequences and hosts without undue experimentation. For example, in selecting a vector, the host must be considered because the vector must replicate in it or be able to integrate into the chromosome. The vector's copy number, the ability to control that copy number, and the expression of any other proteins encoded by the vector, such as antibiotic markers, should also be considered. In selecting an expression control sequence, a variety of factors should also be
20 considered. These include, for example, the relative strength of the sequence, its controllability, and its compatibility with the nucleotide sequence encoding the polypeptide, particularly as regards potential secondary structures. Hosts should be selected by consideration of their compatibility with the chosen vector, the toxicity of the product coded for by the nucleotide sequence, their secretion characteristics, their ability to fold the polypeptide correctly, their
25 fermentation or culture requirements, and the ease of purification of the products coded for by the nucleotide sequence.

The recombinant vector may be an autonomously replicating vector, i.e. a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g. a plasmid. Alternatively, the vector is one which, when introduced
30 into a host cell, is integrated into the host cell genome and replicated together with the chromosome(s) into which it has been integrated.

The vector is preferably an expression vector, in which the nucleotide sequence encoding the polypeptide of the invention is operably linked to additional segments required for transcription of the nucleotide sequence. The vector is typically derived from plasmid or viral
35 DNA. A number of suitable expression vectors for expression in the host cells mentioned herein

003289"69524950

5 are commercially available or described in the literature. Useful expression vectors for eukaryotic hosts, include, for example, vectors comprising expression control sequences from SV40, bovine papilloma virus, adenovirus and cytomegalovirus. Specific vectors are, e.g., pCDNA3.1(+)\Hyg (Invitrogen, Carlsbad, CA, USA) and pCI-neo (Stratagene, La Jola, CA, USA). Useful expression vectors for bacterial hosts include known bacterial plasmids, such as
10 plasmids from *E. coli*, including pBR322, pET3a and pET12a (both from Novagen Inc., WI, USA), wider host range plasmids, such as RP4, phage DNAs, e.g., the numerous derivatives of phage lambda, e.g. , NM989, and other DNA phages, such as M13 and filamentous single stranded DNA phages. Useful expression vectors for yeast cells include the 2 μ plasmid and derivatives thereof, the POT1 vector (US 4,931,373), the pJSO37 vector described in (Okkels,
15 Ann. New York Acad. Sci. 782, 202-207, 1996) and pPICZ A, B or C (Invitrogen). Useful vectors for insect cells include pVL941, pBG311 (Cate et al., "Isolation of the Bovine and Human Genes for Mullerian Inhibiting Substance And Expression of the Human Gene In Animal Cells", Cell, 45, pp. 685-98 (1986), pBluebac 4.5 and pMelbac (both available from Invitrogen).

20 Other vectors for use in this invention include those that allow the nucleotide sequence encoding the polypeptide variant to be amplified in copy number. Such amplifiable vectors are well known in the art. They include, for example, vectors able to be amplified by DHFR amplification (see, e.g., Kaufman, U.S. Pat. No. 4,470,461, Kaufman and Sharp, "Construction Of A Modular Dihydrofolate Reductase cDNA Gene: Analysis Of Signals Utilized For Efficient Expression", Mol. Cell. Biol., 2, pp. 1304-19 (1982)) and glutamine synthetase
25 ("GS") amplification (see, e.g., US 5,122,464 and EP 338,841).

The recombinant vector may further comprise a DNA sequence enabling the vector to replicate in the host cell in question. An example of such a sequence (when the host cell is a mammalian cell) is the SV40 origin of replication. When the host cell is a yeast cell, suitable sequences enabling the vector to replicate are the yeast plasmid 2 μ replication genes REP 1-3
30 and origin of replication.

The vector may also comprise a selectable marker, e.g. a gene the product of which complements a defect in the host cell, such as the gene coding for dihydrofolate reductase (DHFR) or the *Schizosaccharomyces pombe* TPI gene (described by P.R. Russell, Gene 40, 1985, pp. 125-130), or one which confers resistance to a drug, e.g. ampicillin, kanamycin,

5 tetracyclin, chloramphenicol, neomycin, hygromycin or methotrexate. For filamentous fungi, selectable markers include amdS, pyrG, arcB, niaD, sC.

The term "control sequences" is defined herein to include all components, which are necessary or advantageous for the expression of the polypeptide of the invention. Each control sequence may be native or foreign to the nucleic acid sequence encoding the polypeptide.

10 Such control sequences include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, promoter, enhancer or upstream activating sequence, signal peptide sequence, and transcription terminator. At a minimum, the control sequences include a promoter.

A wide variety of expression control sequences may be used in the present invention. Such useful expression control sequences include the expression control sequences associated with structural genes of the foregoing expression vectors as well as any sequence known to control the expression of genes of prokaryotic or eukaryotic cells or their viruses, and various combinations thereof.

Examples of suitable control sequences for directing transcription in mammalian cells include the early and late promoters of SV40 and adenovirus, e.g. the adenovirus 2 major late promoter, the MT-1 (metallothionein gene) promoter, the human cytomegalovirus immediate-early gene promoter (CMV), the human elongation factor 1 α (EF-1 α) promoter, the *Drosophila* minimal heat shock protein 70 promoter, the Rous Sarcoma Virus (RSV) promoter, the human ubiquitin C (UbC) promoter, the human growth hormone terminator, SV40 or adenovirus Elb region polyadenylation signals and the Kozak consensus sequence (Kozak, M. *J Mol Biol* 1987 Aug 20;196(4):947-50).

In order to improve expression in mammalian cells a synthetic intron may be inserted in the 5' untranslated region of the nucleotide sequence encoding the polypeptide of interest. An example of a synthetic intron is the synthetic intron from the plasmid pCI-Neo (available from Promega Corporation, WI, USA).

30 Examples of suitable control sequences for directing transcription in insect cells include the polyhedrin promoter, the P10 promoter, the *Autographa californica* polyhedrosis virus basic protein promoter, the baculovirus immediate early gene 1 promoter and the baculovirus 39K delayed-early gene promoter, and the SV40 polyadenylation sequence.

Examples of suitable control sequences for use in yeast host cells include the promoters of the yeast α -mating system, the yeast triose phosphate isomerase (TPI) promoter,

5 promoters from yeast glycolytic genes or alcohol dehydrogenase genes, the ADH2-4c promoter and the inducible GAL promoter.

Examples of suitable control sequences for use in filamentous fungal host cells include the ADH3 promoter and terminator, a promoter derived from the genes encoding *Aspergillus oryzae* TAKA amylase triose phosphate isomerase or alkaline protease, an *A. niger* 10 α -amylase, *A. niger* or *A. nidulans* glucoamylase, *A. nidulans* acetamidase, *Rhizomucor miehei* aspartic proteinase or lipase, the TPI1 terminator and the ADH3 terminator.

Examples of suitable control sequences for use in bacterial host cells include promoters of the *lac* system, the *trp* system, the *TAC* or *TRC* system and the major promoter regions of phage lambda.

15 The nucleotide sequence of the invention encoding an interferon β polypeptide, whether prepared by site-directed mutagenesis, synthesis or other methods, may or may not also include a nucleotide sequence that encode a signal peptide. The signal peptide is present when the polypeptide is to be secreted from the cells in which it is expressed. Such signal peptide, if present, should be one recognized by the cell chosen for expression of the polypeptide. The 20 signal peptide may be homologous (e.g. be that normally associated with human interferon β) or heterologous (i.e. originating from another source than human interferon β) to the polypeptide or may be homologous or heterologous to the host cell, i.e. be a signal peptide normally expressed from the host cell or one which is not normally expressed from the host cell. Accordingly, the signal peptide may be prokaryotic, e.g. derived from a bacterium such as *E. coli*, or eukaryotic, 25 e.g. derived from a mammalian, or insect or yeast cell.

The presence or absence of a signal peptide will, e.g., depend on the expression host cell used for the production of the polypeptide, the protein to be expressed (whether it is an intracellular or extracellular protein) and whether it is desirable to obtain secretion. For use in filamentous fungi, the signal peptide may conveniently be derived from a gene encoding an 30 *Aspergillus* sp. amylase or glucoamylase, a gene encoding a *Rhizomucor miehei* lipase or protease or a *Humicola lanuginosa* lipase. The signal peptide is preferably derived from a gene encoding *A. oryzae* TAKA amylase, *A. niger* neutral α -amylase, *A. niger* acid-stable amylase, or *A. niger* glucoamylase. For use in insect cells, the signal peptide may conveniently be derived from an insect gene (cf. WO 90/05783), such as the lepidopteran *Manduca sexta* adipokinetic 35 hormone precursor, (cf. US 5,023,328), the honeybee melittin (Invitrogen), ecdysteroid

005280" 6254960

5 UDPglucosyltransferase (egt) (Murphy et al., Protein Expression and Purification 4, 349-357 (1993) or human pancreatic lipase (hpl) (Methods in Enzymology 284, pp. 262-272, 1997). A preferred signal peptide for use in mammalian cells is that of human interferon β apparent from the examples hereinafter or the murine Ig kappa light chain signal peptide (Coloma, M (1992) J. Imm. Methods 152:89-104). For use in yeast cells suitable signal peptides have been found to be
10 the α -factor signal peptide from *S. cerevisiae*. (cf. US 4,870,008), the signal peptide of mouse salivary amylase (cf. O. Hagenbuchle et al., Nature 289, 1981, pp. 643-646), a modified carboxypeptidase signal peptide (cf. L.A. Valls et al., Cell 48, 1987, pp. 887-897), the yeast BAR1 signal peptide (cf. WO 87/02670), and the yeast aspartic protease 3 (YAP3) signal peptide (cf. M. Egel-Mitani et al., Yeast 6, 1990, pp. 127-137).

15 Any suitable host may be used to produce the interferon β polypeptide, including bacteria, fungi (including yeasts), plant, insect, mammal, or other appropriate animal cells or cell lines, as well as transgenic animals or plants. Examples of bacterial host cells include grampositive bacteria such as strains of *Bacillus*, e.g. *B. brevis* or *B. subtilis*, *Pseudomonas* or *Streptomyces*, or gramnegative bacteria, such as strains of *E. coli*. The introduction of a vector
20 into a bacterial host cell may, for instance, be effected by protoplast transformation (see, e.g., Chang and Cohen, 1979, *Molecular General Genetics* 168: 111-115), using competent cells (see, e.g., Young and Spizizin, 1961, *Journal of Bacteriology* 81: 823-829, or Dubnau and Davidoff-Abelson, 1971, *Journal of Molecular Biology* 56: 209-221), electroporation (see, e.g., Shigekawa and Dower, 1988, *Biotechniques* 6: 742-751), or conjugation (see, e.g., Koehler and Thorne,
25 1987, *Journal of Bacteriology* 169: 5771-5278).

 Examples of suitable filamentous fungal host cells include strains of *Aspergillus*, e.g. *A. oryzae*, *A. niger*, or *A. nidulans*, *Fusarium* or *Trichoderma*. Fungal cells may be transformed by a process involving protoplast formation, transformation of the protoplasts, and regeneration of the cell wall in a manner known *per se*. Suitable procedures for transformation of
30 *Aspergillus* host cells are described in EP 238 023 and US 5,679,543. Suitable methods for transforming *Fusarium* species are described by Malardier *et al.*, 1989, *Gene* 78: 147-156 and WO 96/00787. Yeast may be transformed using the procedures described by Becker and Guarente, In Abelson, J.N. and Simon, M.I., editors, *Guide to Yeast Genetics and Molecular Biology, Methods in Enzymology*, Volume 194, pp 182-187, Academic Press, Inc., New York;

- 5 Ito *et al.*, 1983, *Journal of Bacteriology* 153: 163; and Hinnen *et al.*, 1978, *Proceedings of the National Academy of Sciences USA* 75: 1920.

Examples of suitable yeast host cells include strains of *Saccharomyces*, e.g. *S. cerevisiae*, *Schizosaccharomyces*, *Kluyveromyces*, *Pichia*, such as *P. pastoris* or *P. methanolica*, *Hansenula*, such as *H. Polymorpha* or *Yarrowia*. Methods for transforming yeast cells with
10 heterologous DNA and producing heterologous polypeptides therefrom are disclosed by Clontech Laboratories, Inc, Palo Alto, CA, USA (in the product protocol for the Yeastmaker™ Yeast Transformation System Kit), and by Reeves *et al.*, *FEMS Microbiology Letters* 99 (1992) 193-198, Manivasakam and Schiestl, *Nucleic Acids Research*, 1993, Vol. 21, No. 18, pp. 4414-4415 and Ganeva *et al.*, *FEMS Microbiology Letters* 121 (1994) 159-164.

15 Examples of suitable insect host cells include a *Lepidoptera* cell line, such as *Spodoptera frugiperda* (Sf9 or Sf21) or *Trichoplusia ni* cells (High Five) (US 5,077,214). Transformation of insect cells and production of heterologous polypeptides therein may be performed as described by Invitrogen.

Examples of suitable mammalian host cells include Chinese hamster ovary (CHO)
20 cell lines, (e.g. CHO-K1; ATCC CCL-61), Green Monkey cell lines (COS) (e.g. COS 1 (ATCC CRL-1650), COS 7 (ATCC CRL-1651)); mouse cells (e.g. NS/O), Baby Hamster Kidney (BHK) cell lines (e.g. ATCC CRL-1632 or ATCC CCL-10), and human cells (e.g. HEK 293 (ATCC CRL-1573)), as well as plant cells in tissue culture. Additional suitable cell lines are known in the art and available from public depositories such as the American Type Culture Collection,
25 Rockville, Maryland. Also, the mammalian cell, such as a CHO cell, may be modified to express sialyltransferase, e.g. 1,6-sialyltransferase, e.g. as described in US 5,047,335, in order to provide improved glycosylation of the interferon β polypeptide.

Methods for introducing exogenous DNA into mammalian host cells include calcium phosphate-mediated transfection, electroporation, DEAE-dextran mediated transfection,
30 liposome-mediated transfection, viral vectors and the transfection methods described by Life Technologies Ltd, Paisley, UK using Lipofectamin 2000 and Roche Diagnostics Corporation, Indianapolis, USA using FuGENE 6. These methods are well known in the art and e.g. described by Ausbel *et al.* (eds.), 1996, *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, USA. The cultivation of mammalian cells are conducted according to established methods,
35 e.g. as disclosed in (*Animal Cell Biotechnology, Methods and Protocols*, Edited by Nigel

5 Jenkins, 1999, Human Press Inc, Totowa, New Jersey, USA and Harrison MA and Rae IF, General Techniques of Cell Culture, Cambridge University Press 1997).

In the production methods of the present invention, the cells are cultivated in a nutrient medium suitable for production of the polypeptide using methods known in the art. For example, the cell may be cultivated by shake flask cultivation, small-scale or large-scale
10 fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermenters performed in a suitable medium and under conditions allowing the polypeptide to be expressed and/or isolated. The cultivation takes place in a suitable nutrient medium comprising carbon and nitrogen sources and inorganic salts, using procedures known in the art. Suitable media are available from commercial suppliers or may be prepared according to
15 published compositions (e.g., in catalogues of the American Type Culture Collection). If the polypeptide is secreted into the nutrient medium, the polypeptide can be recovered directly from the medium. If the polypeptide is not secreted, it can be recovered from cell lysates.

The resulting polypeptide may be recovered by methods known in the art. For example, the polypeptide may be recovered from the nutrient medium by conventional
20 procedures including, but not limited to, centrifugation, filtration, extraction, spray drying, evaporation, or precipitation.

The polypeptides may be purified by a variety of procedures known in the art including, but not limited to, chromatography (e.g., ion exchange, affinity, hydrophobic, chromatofocusing, and size exclusion), electrophoretic procedures (e.g., preparative isoelectric
25 focusing), differential solubility (e.g., ammonium sulfate precipitation), SDS-PAGE, or extraction (see, e.g., *Protein Purification*, J.-C. Janson and Lars Ryden, editors, VCH Publishers, New York, 1989). Specific methods for purifying polypeptides exhibiting interferon β activity are disclosed in US 4,289,689, US 4,359,389, US 4,172,071, US 4,551,271, US 5,244,655, US 4,485,017, US 4,257,938 and US 4,541,952. A specific purification method is based on
30 immunoaffinity purification (see, e.g., Okamura et al., "Human Fibroblastoid Interferon: Immunosorbent Column Chromatography And N-Terminal Amino Acid Sequence", *Biochem.*, 19, pp. 3831-35 (1980)). Furthermore, purification may be based on the use of IFNAR 1 and/or IFNAR 2, in particular IFNAR 2.

The biological activity of the interferon β polypeptides can be assayed by any
35 suitable method known in the art. Such assays include antibody neutralization of antiviral

5 activity, induction of protein kinase, oligoadenylate 2,5-A synthetase or phosphodiesterase activities, as described in EP 41313 B1. Such assays also include immunomodulatory assays (see, e.g., US 4,753,795), growth inhibition assays, and measurement of binding to cells that express interferon receptors. Specific assays for determining the biological activity of polypeptides or conjugates of the invention are disclosed in the Materials and Methods section
10 hereinafter.

CELL CULTURE OF THE INVENTION

In a further aspect the invention relates to a cell culture comprising a) a host cell transformed with a nucleotide sequence encoding a polypeptide exhibiting interferon β activity, and b) a culture medium comprising said polypeptide produced by expression of said nucleotide
15 sequence in a concentration of at least 800,000 IU/ml of medium, preferably in a concentration in the range of 800,000-3,500,000 IU/ml medium. While the polypeptide exhibiting interferon α activity may be a wild-type interferon β , e.g. human interferon β or a variant thereof (e.g. interferon β 1a or 1b) the polypeptide is preferably an interferon β polypeptide as described herein.

20 In a still further aspect the invention relates to a method of producing an interferon β polypeptide as described herein, the method comprising:

(a) culturing a cell expressing an interferon β polypeptide variant in a culture medium, such that the concentration of the interferon β polypeptide variant in the medium is at least 800,000 IU/ml medium, in particular in the range of between 800,000 and 3,500,000 IU/ml
25 medium; and

(b) recovering the interferon β polypeptide.

OTHER METHODS OF THE INVENTION

In a still further aspect the invention relates to a method reducing immunogenicity and/or of increasing functional *in vivo* half-life and/or serum half-life of an interferon β
30 polypeptide, which method comprises introducing an amino acid residue constituting an attachment group for a first non-polypeptide moiety into a position exposed at the surface of the protein that does not contain such group and/or removing an amino acid residue constituting an

5 attachment group for a first non-polypeptide moiety and subjecting the resulting modified polypeptide to conjugation with the first non-polypeptide moiety.

10 Preferably, the amino acid residue to be introduced and/or removed is as defined in the present application. The non-polypeptide moiety is normally selected from the group consisting of a polymer molecule, a sugar moiety, a lipophilic group and an organic derivatizing agent.

In a still further aspect the invention relates to a method for preparing a conjugate of the invention, wherein the interferon β polypeptide is reacted with the non-polypeptide moiety to which it is to be conjugated under conditions conducive for the conjugation to take place, and the conjugate is recovered.

15 Pharmaceutical composition and uses of a conjugate of the invention

20 The interferon β polypeptide or the conjugate of the invention is administered at a dose approximately paralleling that employed in therapy with human interferon β such as Avonex™, Rebif® and Betaseron®, or a higher dosis. The exact dose to be administered depends on the circumstances. Normally, the dose should be capable of preventing or lessening the severity or spread of the condition or indication being treated. It will be apparent to those of skill in the art that an effective amount of a polypeptide, conjugate or composition of the invention depends, inter alia, upon the disease, the dose, the administration schedule, whether the polypeptide or conjugate or composition is administered alone or in conjunction with other therapeutic agents, the serum half-life of the compositions, and the general health of the patient.

25 The polypeptide or conjugate of the invention can be used "as is" and/or in a salt form thereof. Suitable salts include, but are not limited to, salts with alkali metals or alkaline earth metals, such as sodium, potassium, lithium, calcium and magnesium, as well as e.g. zinc salts. These salts or complexes may be present as a crystalline and/or amorphous structure.

30 The polypeptide or conjugate of the invention is preferably administered in a composition including a pharmaceutically acceptable carrier or excipient. "Pharmaceutically acceptable" means a carrier or excipient that does not cause any untoward effects in patients to whom it is administered. Such pharmaceutically acceptable carriers and excipients are well known in the art.

35 The polypeptide or conjugate of the invention can be formulated into pharmaceutical compositions by well-known methods. Suitable formulations are described in US

5 5,183,746, Remington's Pharmaceutical Sciences by E.W.Martin, 18th edition, A. R. Gennaro, Ed., Mack Publishing Company [1990]; Pharmaceutical Formulation Development of Peptides and Proteins, S. Frokjaer and L. Hovgaard, Eds., Taylor & Francis [2000]; and Handbook of Pharmaceutical Excipients, 3rd edition, A. Kibbe, Ed., Pharmaceutical Press [2000]).

10 The pharmaceutical composition of the polypeptide or conjugate of the invention may be formulated in a variety of forms, including liquid, gel, lyophilized, pulmonary dispersion, or any other suitable form, e.g. as a compressed solid. The preferred form will depend upon the particular indication being treated and will be apparent to one of skill in the art.

15 The pharmaceutical composition containing the polypeptide or conjugate of the invention may be administered orally, intravenously, intracerebrally, intramuscularly, intraperitoneally, intradermally, subcutaneously, intranasally, intrapulmonary, by inhalation, or in any other acceptable manner, e.g. using PowderJect or ProLease technology. The preferred mode of administration will depend upon the particular indication being treated and will be apparent to one of skill in the art.

20 Parenterals

25 An example of a pharmaceutical composition is a solution designed for parenteral administration. Although in many cases pharmaceutical solution formulations are provided in liquid form, appropriate for immediate use, such parenteral formulations may also be provided in frozen or in lyophilized form. In the former case, the composition must be thawed prior to use. The latter form is often used to enhance the stability of the active compound contained in the composition under a wider variety of storage conditions, as it is recognized by those skilled in the art that lyophilized preparations are generally more stable than their liquid counterparts. Such lyophilized preparations are reconstituted prior to use by the addition of one or more suitable pharmaceutically acceptable diluents such as sterile water for injection or sterile physiological saline solution.

30 In case of parenterals, they are prepared for storage as lyophilized formulations or aqueous solutions by mixing, as appropriate, the polypeptide having the desired degree of purity with one or more pharmaceutically acceptable carriers, excipients or stabilizers typically employed in the art (all of which are termed "excipients"), for example buffering agents, stabilizing agents, preservatives, isotonifiers, non-ionic detergents, antioxidants and/or other
35 miscellaneous additives.

Sustained release preparations

Suitable examples of sustained-release preparations include semi-permeable matrices of solid hydrophobic polymers containing the polypeptide or conjugate, the matrices having a suitable form such as a film or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate) or poly(vinylalcohol)), polylactides, copolymers of L-glutamic acid and ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the ProLease® technology or Lupron Depot® (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for long periods such as up to or over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated polypeptides remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

Pulmonary delivery

Conjugate formulations suitable for use with a nebulizer, either jet or ultrasonic, will typically comprise the conjugate dissolved in water at a concentration of, e.g., about 0.01 to 25 mg of conjugate per mL of solution, preferably about 0.1 to 10 mg/mL. The formulation may also include a buffer and a simple sugar (e.g., for protein stabilization and regulation of osmotic pressure), and/or human serum albumin ranging in concentration from 0.1 to 10 mg/ml.

Examples of buffers which may be used are sodium acetate, citrate and glycine. Preferably, the buffer will have a composition and molarity suitable to adjust the solution to a pH in the range of 3 to 9. Generally, buffer molarities of from 1 mM to 50 mM are suitable for this purpose. Examples of sugars which can be utilized are lactose, maltose, mannitol, sorbitol, trehalose, and xylose, usually in amounts ranging from 1% to 10% by weight of the formulation.

The nebulizer formulation may also contain a surfactant to reduce or prevent surface induced aggregation of the protein caused by atomization of the solution in forming the aerosol. Various conventional surfactants can be employed, such as polyoxyethylene fatty acid esters and alcohols, and polyoxyethylene sorbitan fatty acid esters. Amounts will generally range between 0.001% and 4% by weight of the formulation. An especially preferred surfactant for purposes of this invention is polyoxyethylene sorbitan monooleate.

Specific formulations and methods of generating suitable dispersions of liquid particles of the invention are described in WO 9420069, US 5915378, US 5960792, US 5957124, US 5934272, US 5915378, US 5855564, US 5826570 and US 5522385 which are hereby incorporated by reference.

Three specific examples of commercially available nebulizers suitable for the practice of this invention are the Ultravent nebulizer, manufactured by Mallinckrodt, Inc., St. Louis, Mo., the Acorn II nebulizer, manufactured by Marquest Medical Products, Englewood, Colorado, and the AERx pulmonary drug delivery system manufactured by Aradigm Corporation, Hayward, California.

Conjugate formulations for use with a metered dose inhaler device will generally comprise a finely divided powder. This powder may be produced by lyophilizing and then milling a liquid conjugate formulation and may also contain a stabilizer such as human serum albumin (HSA). Typically, more than 0.5% (w/w) HSA is added. Additionally, one or more sugars or sugar alcohols may be added to the preparation if necessary. Examples include lactose, maltose, mannitol, sorbitol, sorbitose, trehalose, xylitol, and xylose. The amount added to the formulation can range from about 0.01 to 200% (w/w), preferably from approximately 1 to 50%, of the conjugate present. Such formulations are then lyophilized and milled to the desired particle size.

The properly sized particles are then suspended in a propellant with the aid of a surfactant. The propellant may be any conventional material employed for this purpose, such as a chlorofluorocarbon, a hydrochlorofluorocarbon, a hydrofluorocarbon, or a hydrocarbon, including trichlorofluoromethane, dichlorodifluoromethane, dichlorotetrafluoroethanol, and 1,1,1,2-tetrafluoroethane, or combinations thereof. Suitable surfactants include sorbitan trioleate and soya lecithin. Oleic acid may also be useful as a surfactant. This mixture is then loaded into the delivery device. An example of a commercially available metered dose inhaler suitable for

5 use in the present invention is the Ventolin metered dose inhaler, manufactured by Glaxo Inc.,
Research Triangle Park, N.C.

Such conjugate formulations for powder inhalers will comprise a finely divided
dry powder containing conjugate and may also include a bulking agent, such as lactose, sorbitol,
sucrose, or mannitol in amounts which facilitate dispersal of the powder from the device, e.g.,
10 50% to 90% by weight of the formulation. The particles of the powder shall have aerodynamic
properties in the lung corresponding to particles with a density of about 1 g/cm² having a median
diameter less than 10 micrometers, preferably between 0.5 and 5 micrometers, most preferably of
between 1.5 and 3.5 micrometers.

An example of a powder inhaler suitable for use in accordance with the teachings
15 herein is the Spinhaler powder inhaler, manufactured by Fisons Corp., Bedford, Mass.

The powders for these devices may be generated and/or delivered by methods
disclosed in US 5997848, US 5993783, US 5985248, US 5976574, US 5922354, US 5785049
and US 55654007 which are hereby incorporated by reference.

The pharmaceutical composition containing the conjugate of the invention may be
20 administered by a wide range of mechanical devices designed for pulmonary delivery of
therapeutic products, including but limited to nebulizers, metered dose inhalers, and powder
inhalers, all of which are familiar to those of skill in the art.

Some specific examples of commercially available devices suitable for the
practice of this invention are the Ultravent nebulizer, manufactured by Mallinckrodt, Inc., St.
25 Louis, Missouri; the Acorn II nebulizer, manufactured by Marquest Medical Products,
Englewood, Colorado; the Ventolin metered dose inhaler, manufactured by Glaxo Inc., Research
Triangle Park, North Carolina; the Spinhaler powder inhaler, manufactured by Fisons Corp.,
Bedford, Massachusetts; the "standing cloud" device of Inhale Therapeutic Systems, Inc., San
Carlos, California; the AIR inhaler manufactured by Alkermes, Cambridge, Massachusetts; and
30 the AERx pulmonary drug delivery system manufactured by Aradigm Corporation, Hayward,
California.

The pharmaceutical composition of the invention may be administered in
conjunction with other therapeutic agents. These agents may be incorporated as part of the same
pharmaceutical composition or may be administered separately from the polypeptide or
35 conjugate of the invention, either concurrently or in accordance with any other acceptable

5 treatment schedule. In addition, the polypeptide, conjugate or pharmaceutical composition of the invention may be used as an adjunct to other therapies.

Accordingly, this invention provides compositions and methods for treating most types of viral infections, cancers or tumors (e.g. breast carcinoma, non-small cell lung cancer) or tumour angiogenesis, Chrohn's disease, ulcerative colitis, Guillain-Barré syndrome, glioma,
10 idiopathic pulmonary fibrosis, abnormal cell growth, or for immunomodulation in any suitable animal, preferably mammal, and in particular human. In particular the polypeptide, conjugate or composition of the invention may be used for the treatment of multiple sclerosis (MS), such as any of the generally recognized four types of MS (benign, relapsing remitting MS (RRMS), primary progressive MS (PPMS) and secondary progressive MS (SPMS)) and for
15 monosymptomatic MS), hepatitis, or a herpes infection (the latter treatment optionally being combined with a treatment with IL-10).

In a further aspect the invention relates to a method of treating a mammal having circulating antibodies against interferon β 1a, such as AvonexTM or Rebif®, or 1b, such as Betaseron®, which method comprises administering a compound which has the bioactivity of
20 interferon β and which has a reduced or no reaction with said antibodies. The compound is administered in an effective amount. The compound is preferably a conjugate as described herein and the mammal is preferably a human being. The mammals to be treated may suffer from any of the diseases listed above for which interferon β is a useful treatment. In particular, this aspect of the invention is of interest for the treatment of multiple sclerosis (any of the types listed above)
25 or cancer. Furthermore, the invention relates to a method of making a pharmaceutical product for use in treatment of mammals having circulating antibodies against interferon β 1a, such as AvonexTM or Rebif®, or 1b, such as Betaseron®, wherein a compound which has the bioactivity of interferon β and which does not react with such is formulated into an injectable or otherwise suitable formulation. The term "circulating antibodies" is intended to indicate antibodies, in
30 particular neutralizing antibodies, formed in a mammal in response to having been treated with any of the commercially available interferon β preparations (Rebif®, Betaseron®, AvonexTM).

In a further aspect the invention relates to a method of treating a patient in need of treatment with a pharmaceutical composition with at least some of the therapeutically beneficial properties of interferon β comprising administering a composition comprising a compound with
35 at least part of the therapeutically beneficial activity of interferon β , said treatment having

5 reduced or removed adverse psychological effects as compared to treatment with interferon β , wherein said compound is a non-naturally occurring conjugate of a polypeptide with interferon β activity and a non-polypeptide moiety, in particular a conjugate according to the present invention.

10 In a still further aspect the invention relates to a pharmaceutical composition for the treatment of a patient in need of treatment with a compound having at least part of the therapeutically beneficial properties of interferon β , said composition comprising a compound which is a non-naturally occurring conjugate of interferon β and a non-polypeptide moiety, said treatment further giving rise to fewer adverse psychological effects than treatment with interferon β . The conjugate is preferably a conjugate of the invention.

15 Also contemplated is use of a nucleotide sequence encoding a polypeptide of the invention in gene therapy applications. In particular, it may be of interest to use a nucleotide sequence encoding a polypeptide as described in the section above entitled "Glycosylated Polypeptides of the Invention modified to incorporate additional glycosylation sites". The glycosylation of the polypeptides is thus achieved during the course of the gene therapy, i.e. after
20 expression of the nucleotide sequence in the human body.

Gene therapy applications contemplated include treatment of those diseases in which the polypeptide is expected to provide an effective therapy due to its antiviral activity, e.g., viral diseases, including hepatitis such as hepatitis C, and particularly HPV, or other infectious diseases that are responsive to interferon β or infectious agents sensitive to interferon
25 β . Furthermore, the conjugate or polypeptide of the invention may be used in the treatment of chronic inflammatory demyelinating polyradiculoneuropathy, and of severe necrotising cutaneous lesions. Also, gene therapy in connection with the treatment of any MS type is contemplated. Similarly, this invention contemplates gene therapy applications for immunomodulation, as well as in the treatment of those diseases in which interferon β is
30 expected to provide an effective therapy due to its antiproliferative activity, e.g., tumors and cancers, or other conditions characterized by undesired cell proliferation, such as restenosis. A further description of such gene therapy is provided in WO 95/25170.

Local delivery of interferon β using gene therapy may provide the therapeutic agent to the target area while avoiding potential toxicity problems associated with non-specific
35 administration.

Both *in vitro* and *in vivo* gene therapy methodologies are contemplated.

Several methods for transferring potentially therapeutic genes to defined cell populations are known. For further reference see, e.g., Mulligan, "The Basic Science Of Gene Therapy", Science, 260, pp. 926-31 (1993). These methods include:

Direct gene transfer, e.g., as disclosed by Wolff et al., "Direct Gene transfer Into Mouse Muscle In vivo", Science 247, pp. 1465-68 (1990);

Liposome-mediated DNA transfer, e.g., as disclosed by Caplen et al., "Liposome-mediated CFTR Gene Transfer to the Nasal Epithelium Of Patients With Cystic Fibrosis" Nature Med., 3, pp. 39-46 (1995); Crystal, "The Gene As A Drug", Nature Med., 1, pp.- 15-17 (1995); Gao and Huang, "A Novel Cationic Liposome Reagent For Efficient Transfection of Mammalian Cells", Biochem.Biophys Res. Comm., 179, pp. 280-85 (1991);

Retrovirus-mediated DNA transfer, e.g., as disclosed by Kay et al., "In vivo Gene Therapy of Hemophilia B: Sustained Partial Correction In Factor IX-Deficient Dogs", Science, 262, pp. 117-19 (1993); Anderson, "Human Gene Therapy", Science, 256, pp.808-13(1992);

DNA Virus-mediated DNA transfer. Such DNA viruses include adenoviruses (preferably Ad-2 or Ad-5 based vectors), herpes viruses (preferably herpes simplex virus based vectors), and parvoviruses (preferably "defective" or non-autonomous parvovirus based vectors, more preferably adeno-associated virus based vectors, most preferably AAV-2 based vectors). See, e.g., Ali et al., "The Use Of DNA Viruses as Vectors for Gene Therapy", Gene Therapy, 1, pp. 367-84 (1994); US 4,797,368, and US 5,139,941.

The invention is further described in the following examples. The examples should not, in any manner, be understood as limiting the generality of the present specification and claims.

MATERIALS AND METHODS

Materials

HeLa cells – (available from American Type Culture Collection (ATCC))
ISRE-Luc (Stratagene, La Jolla USA)
pCDNA 3.1/hygro (Invitrogen, Carlsbad USA)
pGL3 basic vector (Promega)
Human genomic DNA (CloneTech, USA)

- 5 DMEM medium: Dulbecco's Modified Eagle Media (DMEM), 10% fetal bovine serum
(available from Life Technologies A/S, Copenhagen, Denmark)

ASSAYS

Interferon Assay

- 10 It has previously been published that interferon β interacts with and activates
Interferon type I receptors on HeLa cells. Consequently, transcription is activated at promoters
containing an Interferon Stimulated Response Element (ISRE). It is thus possible to screen for
agonists of interferon receptors by use of an ISRE coupled luciferase reporter gene (ISRE-luc)
placed in HeLa cells.

Primary Assay

- 15 HeLa cells are co-transfected with ISRE-Luc and pCDNA 3.1/hygro and foci (cell
clones) are created by selection in DMEM media containing Hygromycin B. Cell clones are
screened for luciferase activity in the presence or absence of interferon β . Those clones showing
the highest ratio of stimulated to unstimulated luciferase activity are used in further assays.
- 20 To screen muteins, 15,000 cells/well are seeded in 96 well culture plates and
incubated overnight in DMEM media. The next day muteins as well as a known standard are
added to the cells in various concentrations. The plates are incubated for 6 hours at 37 C in a
5% CO₂ air atmosphere LucLite substrate (Packard Bioscience, Groningen The Netherlands) is
subsequently added to each well. Plates are sealed and luminescence measured on a TopCount
- 25 luminometer (Packard) in SPC (single photon counting) mode. Each individual plate contains
wells incubated with interferon β as a stimulated control and other wells containing normal
media as an unstimulated control. The ratio between stimulated and unstimulated luciferase
activity serves as an internal standard for both mutein activity and experiment-to-experiment
variation.

Secondary Assay

- 30 Currently, there are 18 non-allelic interferon α genes and one interferon β gene.
These proteins exhibit overlapping activities and thus it is critical to ensure that muteins retain
the selectivity and specificity of interferon β .

- 35 The β -R1 gene is activated by interferon β but not by other interferons. The
transcription of β -R1 thus serves as a second marker of interferon β activation and is used to

5 ensure that muteins retain interferon β activity. A 300 bp promoter fragment of β -R1 shown to drive interferon sensitive transcription (Rani. M.R. et al (1996) *JBC* **271** 22878-22884) was isolated by PCR from human genomic DNA and inserted into the pGL3 basic vector (Promega). The resulting β -R1:luciferase gene is used in assays similar to the primary assay described above. In astrocytoma cells, the resulting β -R1:luciferase gene has been described to show 250
10 fold higher sensitivity to interferon β than to interferon α (Rani et al. *op cit*).

ELISA assay

The concentration of IFN- β is quantitated by use of a commercial sandwich immunoassay (PBL Biomedical Laboratories, New Brunswick, NJ, USA). The kit is based on an ELISA with monoclonal mouse anti-IFN- β antibodies for catching and detection of IFN- β in test
15 samples. The detecting antibody is conjugated to biotin.

Tests samples and recombinant human IFN- β standard are added in 0.1 mL in concentrations from 10-0.25 ng/mL to microtiter plates, precoated with catching antibody. The plates are incubated at RT for 1 hr. Samples and standard are diluted in kit dilution buffer.

The plates are washed in the kit buffer and incubated with the biotinylated
20 detecting antibody in 0.1 mL for 1 hr at RT. After another wash the streptavidin-horseradishperoxidase conjugate is added in 0.1 mL and incubated for 1 hr at RT.

The reaction is visualised by addition of 0.1 mL Tetramethylbenzidine (TMB) substrate chromogen. The plates are incubated for 15 minutes in the dark at RT and the reaction is stopped by addition of stop solution. The absorbance is read at 450nm using an ELISA reader.

Receptor binding assay

25 The receptor binding capability of a polypeptide or conjugate of the invention can be determined using the assay described in WO 95/25170 entitled "Analysis Of IFN- β (Phe₁₀₁) For Receptor Binding"(which is based on Daudi or A549 cells). Soluble domains of IFNAR1 and IFNAR2 can be obtained essentially as described by Arduini et al, Protein Science, 1999,
30 vol. 8, 1867-1877 or as described in Example 9 herein.

Alternatively, the receptor binding capability is determined using a crosslinking agent such as disuccinimidyl suberate (DSS) available from Pierce, Rockford, IL, USA as follows:

The polypeptide or conjugate is incubated with soluble IFNAR-2 receptor in the
35 presence or absence of DSS in accordance with the manufacturer's instructions. Samples are

5 separated by SDS-PAGE, and a western blot using anti-interferon β or anti-IFNAR2 antibodies is performed. The presence of a functional interferon β polypeptide/conjugate: receptor interaction is apparent by an increase in the molecular size of receptor and interferon β in the presence of DSS.

10 Furthermore, a crosslinking assay using a polypeptide or conjugate of the invention and both receptor subunits (IFNAR-1 and IFNAR-2) can establish Interferon receptor 1 binding ability. In this connection it has been published that IFNAR-1 binds only after an interferon β : IFNAR-2 complex is formed (Mogensen et al., Journal of Interferon and Cytokine Research, 19:1069-1098, 1999).

In vitro immunogenicity tests of interferon β conjugates

15 Reduced immunogenicity of a conjugate or polypeptide of the invention is determined by use of an ELISA method measuring the immunoreactivity of the conjugate or polypeptide relative to a reference molecule or preparation. The reference molecule or preparation is normally a recombinant human interferon β preparation such as Avonex, Rebif or Betaseron, or another recombinant human interferon β preparation produced by a method
20 equivalent to the way these products are made. The ELISA method is based on antibodies from patients treated with one of these recombinant interferon β preparations. The immunogenicity is considered to be reduced when the conjugate or polypeptide of the invention has a statistically significant lower response in the assay than the reference molecule or preparation.

25 Another method of determining immunogenicity is by use of sera from patients treated with interferon beta (i.e. any commercial interferon β product) in an analogous manner to that described by Ross et al. J. Clin Invest. 95, 1974-78, 1995. In the antiviral neutralisation bioassay reduced immunogenicity results in reduced inhibition of a conjugate of the invention by patient sera compared to a wt IFN-beta reference molecule. Furthermore, in the biochemical IFN binding assay a less immunogenic conjugate is expected to bind to patient IgG to a lesser extent
30 than reference IFN-beta molecules.

For the neutralisation assay, the reference and conjugate molecules are added in a concentration that produces approximately 80% virus protection in the antiviral neutralisation bioassay. The IFN- β proteins are mixed with patient sera in various dilutions (starting at 1:20).

5

Antiviral activity

The antiviral bioassay is performed using A549 cells (CCL 185, American tissue culture collection) and Encephalomyocarditis (EMC) virus (VR-129B, American tissue culture collection).

10 The cells are seeded in 96 well tissue culture plates at a concentration of 10,000 cells/well and incubated at 37°C in a 5% CO₂ air atmosphere. A polypeptide or conjugate of the invention is added in concentrations from 100-0.0001 IU/mL in a total of 100µl DMEM medium containing fetal calf serum and antibiotics.

15 After 24 hours the medium is removed and 0.1 mL fresh medium containing EMC virus is added to each well. The EMC virus is added in a concentration that causes 100% cell death in IFN-β free cell cultures after 24 hours.

20 After another 24 hrs, the antiviral effect of the polypeptide or conjugate is measured using the WST-1 assay. 0.01 mL WST-1 (WST-1 cell proliferation agent, Roche Diagnostics GmbH, Mannheim, Germany) is added to 0.1 mL culture and incubated for ½-2 hours at 37°C in a 5% CO₂ air atmosphere. The cleavage of the tetrazolium salt WST-1 by mitochondrial dehydrogenases in viable cells results in the formation of formazan that is quantified by measuring the absorbance at 450 nm.

Neutralisation of activity in Interferon Stimulated Response Element (ISRE) assay

25 The interferon β neutralising effect of anti-interferon β sera are analysed using the ISRE-Luciferase activity assay.

30 Sera from interferon β treated patients or from immunised animals are used. Sera are added either in a fixed concentration (dilution 1:20-1:500 (pt sera) or 20-600 ng/mL (animal sera)) or in five-fold serial dilutions of sera starting at 1/20 (pt sera) or 600 ng/mL (animal sera). Interferon β is added either in five fold-dilutions starting at 25.000 IU/mL or in a fixed concentration (0.1-10 IU/mL) in a total volume of 80µl DMEM medium + 10% FCS. The sera are incubated for 1 hr. at 37°C with IFN-β.

35 The samples are then transferred to 96 well tissue culture plates containing HeLa cells transfected with ISRE-Luc grown from 24 hrs before (15,000 cells/well) in DMEM media. The cultures are incubated for 6 hours at 37°C in a 5% CO₂ air atmosphere. LucLite substrate (Packard Bioscience, Groningen, The Netherlands) is subsequently added to each well. Plates

are sealed and luminescence measured on a TopCount luminometer (Packard) in SPC (single photon counting) mode.

When interferon β samples are titrated in the presence of a fixed amount of serum, the neutralising effect was defined as fold inhibition (FI) quantified as $EC_{50}(w. \text{ serum})/EC_{50}(w/o \text{ serum})$. The reduction of antibody neutralisation of interferon β variant proteins is defined as

$$(1 - \frac{FI \text{ variant}}{FI \text{ wt}}) \times 100\%$$

Biological half-life measurement of a PEG - interferon β conjugate

Measurement of biological half-life can be carried out in a number of ways described in the literature. One method is described by Munafo et al (European Journal of Neurology 1998, vol 5 No2 p 187-193), who used an ELISA method to detect serum levels of interferon β after subcutaneous and intramuscular administration of interferon β .

The rapid decrease of interferon β serum concentrations after i.v. administration has made it important to evaluate biological responses to interferon β treatment. However it is contemplated that the conjugates of the present invention will have prolonged serum half lives also after i.v. administration making it possible to measure by e.g. an ELISA method or by the primary screening assay.

Different pharmacodynamic markers (e.g. serum neuplerin and beta2 microglobulin) have also been studied (Clin Drug Invest (1999) 18(1):27-34). These can equally well be used to evaluate prolonged biological effect. These experiments may also be carried out in suitable animal species, e.g. rats.

Assays to assess the biological effects of interferon β such as antiviral, antiproliferative and immunomodulatory effects (as described in e.g. Annals of Neurology 1995 vol 37 No 1 p 7-15) can be used together with the primary and secondary screening assays described herein to evaluate the biological efficacy of the conjugate in comparison to wild type interferon β .

Finally an animal model such as the commonly used experimental autoimmune encephalomyelitis (EAE) model can be used to establish efficacy of a conjugate or polypeptide of the invention. In the EAE model immunization with myelin or myelin derived proteins elicits a disease mimicking the majority of the inflammatory and neurologic features of multiple

5 sclerosis in humans. EAE has been used in mice, rats, rabbits, and marmosets {Cannella, Hoban, et al. 1998 ID: 695} {Zaprianova, Deleva, et al. 1997 ID: 699} {Hassouna, Galeano, et al. 1983 ID: 731} {Genain & Hauser 1997 ID: 703}. Other models include Theiler's murine encephalomyelitis virus (TMEV) model {Murray, McGavern, et al. 1998 ID: 694}. will be used to establish efficacy of the interferon β conjugate.

10 PEGylation in microtiter plates of a tagged polypeptide with interferon β activity
The method comprises

Expressing the interferon β polypeptide with a suitable tag, e.g. any of the tags exemplified in the general description above.

Transferring culture broth to one or more wells in a microtiter plate capable of
15 immobilising the tagged polypeptide. When the tag is His-His-His-His-His (Casey et al, J. Immunol. Meth., 179, 105 (1995)), a Ni-NTA HisSorb microtiter plate commercially available from QiaGen can be used.

After allowing for immobilising the tagged polypeptide to the microtiter plate, the wells are washed in a buffer suitable for binding and subsequent PEGylation.

20 Incubating the wells with the activated PEG of choice. As an example, M-SPA-5000 from Shearwater Polymers is used. The molar ratio of activated PEG to polypeptide has to be optimised, but will typically be greater than 10:1 more typically greater than 100:1.

After a suitable reaction time at ambient temperature, typically around 1 hour, the reaction is stopped by removal of the activated PEG solution. The conjugated protein is eluted
25 from the plate by incubation with a suitable buffer. Suitable elution buffers may contain Imidazole, excess NTA or another chelating compound.

The conjugated protein is assayed for biological activity and immunogenicity as appropriate.

This tag may optionally be cleaved off using a method known in the art, e.g. using
30 diaminopeptidase and the Gln in pos -1 will be converted to pyroglutamyl with GCT (glutamylcyclotransferase) and finally cleaved off with PGAP (pyro-glutamyl-aminopeptidase) giving the native protein. The process involves several steps of metal chelate affinity chromatography. Alternatively, the tagged polypeptide may be conjugated.

5 PEGylation of a receptor-bound interferon β polypeptide

In order to optimize PEGylation of an interferon β polypeptide in a manner excluding PEGylation of lysines involved in receptor recognition, the following method has been developed: The soluble domains of IFNAR1 and IFNAR2 are obtained essentially as described in Arduini et al, Protein Science (1999), vol 8: 1867-1877 or as described in Example 9.

10 A ternary complex consisting of an interferon β polypeptide, a soluble domain of IFNAR1 and a soluble domain of IFNAR2 in a 1:1:1 stoichiometry is formed in a PBS buffer at pH 7-9. The concentration of Interferon β polypeptide is approximately 20 ug/ml or 1 uM and the receptors are present at equimolar concentration.

M-SPA-5000 from Shearwater Polymers, Inc is added at 3 different concentration
15 levels corresponding to 5, 20 or 100 molar excess of interferon β polypeptide. The reaction time is 30 min at RT. After the 30 min reaction period, the pH of the reaction mixture is adjusted to pH 2.0 and the reaction mixture is applied to a Vydac C18 column and eluted with an acetonitrile gradient essentially as described (Utsumi et al, J. Biochem., vol 101, 1199-1208, (1987). Alternatively and more elegantly, an isopropanol gradient can be used.

20 Fractions are analyzed using the primary screening assay described herein and active PEGylated interferon- β polypeptide obtained by this method stored at -80°C in PBS, pH 7 containing 1 mg/ml HSA.

Alternatively, to the procedure described above a soluble domain of IFNAR2 is used as the only receptor component to form a binary complex. Furthermore, IFNAR2 may be
25 immobilized on a suitable resin (e.g. Epoxy activated Sepharose 6B) according to the manufactures instructions prior to forming the binary complex. After PEGylation, the PEGylated Interferon- β is eluted with a 0.1 M Glycin, pH 2 buffer and activity measured as described after pH adjustment to neutral.

30 Accessible Surface Area (ASA)

The computer program Access (B. Lee and F.M.Richards, J. Mol.Biol. 55: 379-400 (1971)) version 2 (Copyright (c) 1983 Yale University) are used to compute the accessible surface area (ASA) of the individual atoms in the structure. This method typically uses a probe-size of 1.4\AA and defines the Accessible Surface Area (ASA) as the area formed by the centre of the probe. Prior to this calculation all water molecules and all hydrogen atoms are removed from
35 the coordinate set, as are other atoms not directly related to the protein. Alternative programs are

5 available for computing ASA, e.g. the program WhatIf G.Vriend, J. Mol. Graph. (1990) 8, 52-56,
electronically available at the WWW interface on <http://swift.embl-heidelberg.de/servers2/>
(R.Rodriguez *et.al.* CABIOS (1998) 14, 523-528.) using the option *Accessibility* to calculate the
accessible molecular surface.

Fractional ASA of side chain

10 The fractional ASA of the side chain atoms is computed by division of the sum of
the ASA of the atoms in the side chain with a value representing the ASA of the side chain atoms
of that residue type in an extended ALA-x-ALA tripeptide. See Hubbard, Campbell & Thornton
(1991) J.Mol.Biol.220,507-530. For this example the CA atom is regarded as a part of the side
chain of Glycine residues but not for the remaining residues. The following table indicates the

15 100% ASA standard for the side chain:

Ala	69.23 Å ²
Arg	200.35 Å ²
Asn	106.25 Å ²
Asp	102.06 Å ²
Cys	96.69 Å ²
Gln	140.58 Å ²
Glu	134.61 Å ²
Gly	32.28 Å ²
His	147.00 Å ²
Ile	137.91 Å ²
Leu	140.76 Å ²
Lys	162.50 Å ²
Met	156.08 Å ²
Phe	163.90 Å ²
Pro	119.65 Å ²
Ser	78.16 Å ²
Thr	101.67 Å ²
Trp	210.89 Å ²
Tyr	176.61 Å ²
Val	114.14 Å ²

Determining surface exposed amino acid residues

20 The three-dimensional crystal structure of human interferon beta at 2.2 Å
resolution (Karpusas *et al.* Proc. Nat. Acad. Sci. USA (1997) 94:11813-11818 is available from
the Protein Data Bank (PDB) (Bernstein *et.al.* J. Mol. Biol. (1977) 112 pp. 535) and

5 electronically available via The Research Collaboratory for Structural Bioinformatics PDB at <http://www.pdb.org/> under accession code 1AU1. This crystal structure contain two independent molecules of human interferon beta in this example the A molecule is used.

Surface exposure:

10 Using the WhatIf program as described above the following residues were found to have zero surface accessibility for their side chain atoms (for Gly the accessibility of the CA atom is used): G7, N14, C17, L21, I44, A55, A56, T58, I59, M62, L63, L98, L122, Y125, I129, L133, A142, W143, V146, I150, N153, I157, L160, T161, and L164.

Fractional surface exposure

15 For further analysis it was necessary to remodel the side chains of residues R71, R113, K115, L116, M117 due to steric clashes. The remodelling was done using Modeler 98, MSI INC. Performing fractional ASA calculations using the Access computer program on the remodelled interferon beta molecule (only including the amino acid residues and excluding the N-linked sugar moiety) resulted in the following residues having more than 25% of their side chain exposed to the surface: S2, N4, L5, F8, L9, R11, S12, F15, Q16 Q18, K19, W22, Q23, 20 G26, R27, L28, E29, Y30, L32, K33, R35, M36, N37, D39, E42, K45, Q46, L47, Q48, Q49, Q51, K52, Q64, A68, R71, Q72, D73, S75, S76, G78, N80, E81, T82, E85, N86, A89, Y92, H93, N96, H97, K99, T100, E103, E104, K105, E107, K108, E109, D110, F111, R113, G114, K115, L116, S119, L120, H121, K123, R124, G127, R128, L130, H131, K134, A135, K136, E137, Y138, S139, H140, V148, R152, Y155, N158, G162, Y163, R165, and N166. and the 25 following residues have more than 50% of their side chain exposed to the surface: N4, L5, F8, S12, F15, Q16, K19, W22, G26, R27, E29, Y30, K33, R35, N37, D39, E42, Q46, Q48, Q49, Q51, K52, R71, D73, S75, G78, N80, E81, T82, E85, N86, A89, Y92, H93, K99, T100, E103, E104, E107, K108, D110, F111, L116, K123, R124, G127, H131, K134, E137, V148, Y155, R165, and N166.

30

EXAMPLES

EXAMPLE 1. DESIGN OF AN EXPRESSION CASSETTE FOR EXPRESSION OF INTERFERON β IN MAMMALIAN AND INSECT CELLS

The DNA sequence, GenBank accession number M28622 (shown in SEQ ID NO 1), encompassing a full length cDNA encoding human interferon β with its native signal peptide, 35 was modified in order to facilitate high expression in mammalian cells. First the ATG start codon

context was modified according to the Kozak consensus sequence (Kozak, M. *J Mol Biol* 1987 Aug 20;196(4):947-50), such that there is a perfect match to the consensus sequence upstream of the ATG start codon. Secondly the codons of the native human interferon β was modified by making a bias in the codon usage towards the codons frequently used in highly expressed human genes. Subsequently, certain nucleotides in the sequence were substituted with others in order to introduce recognition sites for DNA restriction endonucleases (this allows for easier modification of the DNA sequence later). Primers were designed such that the gene could be synthesised:

CBProFpr1

5'GGCTAGCGTTTAAACTTAAGCTTCGCCACCATGACCAACAAGTGCCTGCTC
CAGATCGCCCTGCTCCTGT-3', SEQ ID 3,

CBProFpr2

5'ACAACCTGCTCGGCTTCCTGCAGAGGAGTTCGAACTTCCAGTGCCAGAAGC
TCCTGTGGCAGCTGAACGG-3', SEQ ID 4,

CBProFpr3

5'GAACTTCGACATCCCCGAGGAAATCAAGCAGCTGCAGCAGTTCCAGAAGG
AGGACGCCGCTCTGACCATC-3', SEQ ID 5,

CBProFpr4

5'TTCCGCCAGGACTCCAGCTCCACCGGTTGGAACGAGACCATCGTGGAGAAC
CTGCTGGCCAACGTGTACC-3', SEQ ID 6,

CBProFpr5

5'AGGAGAAGCTGGAGAAGGAGGACTTCACCCGCGGCAAGCTGATGAGCTCC
CTGCACCTGAAGCGCTACTA-3', SEQ ID 7,

CBProFpr6

5'GGAGTACAGCCACTGCGCCTGGACCATCGTACGCGTGGAGATCCTGCGCAA
CTTCTACTTCATCAACCGC-3', SEQ ID 8,

CBProFpr9

5'CACCACACTGGACTAGTGGATCCTTATCAGTTGCGCAGGTAGCCGGTCAGG
CGGTTGATGAAGTAGAAGT-3', SEQ ID 9,

CBProFpr10

5'AGGCGCAGTGGCTGTACTCCTTGGCCTTCAGGTAGTGCAGGATGCGGCCAT
AGTAGCGCTTCAGGTGCAG-3', SEQ ID 10,

CBProFpr11

5'CTCCTTCTCCAGCTTCTCCTCCAGCACGGTCTTCAGGTGGTTGATCTGGTGG
TACACGTTGGCCAGCAGG-3', SEQ ID 11,

CBProFpr12

5'GAGCTGGAGTCCTGGCGGAAGATGGCGAAGATGTTCTGCAGCATCTCGTAG
ATGGTCAGAGCGGCGTCCT-3', SEQ ID 12,

CBProFpr13

5'CCTCGGGGATGTCGAAGTTCATCCTGTCCTTCAGGCAGTACTCCAGGCGCC
CGTTCAGCTGCCACAGGAG-3', SEQ ID 13,

5 CBProFpr14
5'CAGGAAGCCGAGCAGGTTGTAGCTCATCGATAGGGCCGTGGTGCTGAAGC
ACAGGAGCAGGGCGATCTGG-3', SEQ ID 14,

The primers were assembled to the synthetic gene by one step PCR using
Platinum *Pfx*- polymerase kit (Life Technologies) and standard three step PCR cycling
10 parameters. The assembled gene was amplified by PCR using the same conditions.

A cDNA encoding a N-terminal extended form of human interferon β was
synthesised using the same PCR conditions as described above but with the primers CBProFpr1
and -14 substituted with the primers:

15 CBProFpr7
5'CTGCTCCAGATCGCCCTGCTCCTGTGCTTCAGCACCACGGCCCTATCGATGA
AGCACCAGCACCAGCATC-3', SEQ ID 15,

CBProFpr8
5'CACTGCTTACTGGCTTATCGAAATTAATACGACTCACTATAGGGAGACCCA
20 AGCTGGCTAGCGTTTAAAC-3', SEQ ID 16,

CBProFpr15
5'CAGGAAGCCGAGCAGGTTGTAGCTCATCTGTTGGTGTTGATGTTGGTGCTG
ATGCTGGTGCTGGTGCTTC-3', SEQ ID 17,

CBProFpr16
25 5'AGCAGGGCGATCTGGAGCAGGCACTTGTTGGTCATGGTGGCGAAGCTTAAG
TTTAAACGCTAGCCAGCTT-3', SEQ ID 18,
in order to incorporate a purification TAG in the interferon β molecule.

The synthesised genes were cloned into pcDNA3.1/Hygro (Invitrogen) between
the *Hind*III site at the 5' end and the *Bam*HI at the 3', resulting in pCBProF1 and pCBProF2.

30 The synthetic intron from pCI-Neo (Promega) was amplified using standard PCR
conditions as described above and the primers:

CBProFpr37 5'-CCGTCAGATCCTAGGCTAGCTTATTGCGGTAGTTTATCAC-3', SEQ ID
19,

CBProFpr38 5'-GAGCTCGGTACCAAGCTTTTAAGAGCTGTAAT-3', SEQ ID 20,

35 resulting in a 332 bp PCR fragment which was cut with *Nhe*I and *Hind*III and inserted in the
5'UTR of the plasmids pCBProF1 and pCBProF2 resulting in pCBProF4 and pCBProF5.

Codons for individual amino acids were changed by amplifying relevant regions
of the coding region by PCR in such a way that the PCR introduced changes in the sequence can
be introduced in the expression plasmids by classical cloning techniques. E.g. the primers:

40 Lys45arg-5'primer (*Nar*I/*Kas*I):
5'GCTGAACGGGCGCCTGGAGTACTGCCTGAAGGACAGGATGAACTTCGACATCCCC
GAGGAAATCCGCCAGCTGCAGC-3', SEQ ID 21,

5 Lys45mut-3' primer (BsiWI): 5'TCTCCACGCGTACGATGGTCCAGGCGCAGTGGCTG-3',
SEQ ID 22,
were used to introduce a K45R substitution in the PCR-fragment spanning the region from
position 1055 to 1243 in pCBProF1. Both the PCR fragment and pCBProF1 was cut with NarI
and BsiWI which are both unique. The PCR fragment and the vector backbone of pCBProF1 are
10 purified and ligated resulting in substitution of the Lys45 codon AAG with the Arg codon CGC
in pCBProF1.

Furthermore, SOE (sequence overhang extension) PCR was used for introduction
of amino acid substitutions. In the SOE-PCR both the N-terminal part and the C-terminal part of
the INFB molecule were first amplified in individual primary PCRs.

15 For these primary PCRs the central complementary primers were synthesised such
that the codon(s) for the amino acid(s) to be substituted is/are changed to the desired codon(s).
The terminal primers were standard primers defining the N- and C-terminal of the INFβ
molecule respectively. Further the terminal primers provided a restriction enzyme site enabling
subsequent cloning of the full-length PCR product. Thus, the central (nonsense) primer and the
20 N-terminal (sense) primer were used to amplify the N-terminal part of the INFβ coding region in
one of the primary PCRs and equivalently for the C-terminal part. Once amplified the N- and C-
terminal parts are assembled into the full-length product in a secondary PCR and cloned into a
modified version of pCDNA3.1/Hygro as described above. For instance, the following primers
were used to introduce the mutations for the substitutions F111N and R113T:

25 CBProFprimer9(Sense):
CACCACACTGGACTAGTGGATCCTTATCAGTTGCGCAGGTAGCCGGTCAGGC
GGTTG ATG AAGTAGAAGT (SEQ ID NO 23),

30 CBProFprimer231(Antisense):
CATCAGCTTGCCGGTGGTGTTCCTCCTTC (SEQ ID NO 24),

CBProFprimer230 (Sense):
GAAGGAGGACAACACCACCGCAAGCTGATG (SEQ ID NO 25),

35 CBProFprimer42 (Antisense):
CACACTGGACTAGTAAGCTTTTATCAGTTGCGCAGGTAGC (SEQ ID NO 26),

Furthermore, in cases where the introduced mutation(s) were sufficiently close to
a unique restriction endo-nuclease site in the expression plasmid variant genes were constructed

5 using construction procedure encompassing a single PCR step and a subsequent cloning. For instance, the substitution K19R was introduced by use of the PCR primer:

CBProFpr58:

GAGGAGTTCGAACTTCCAGTGCCAGCGCCTCCTGTGGCAGCTGAACG (SEQ ID

NO 27), and CBProFprimer9:

10 The PCR product was subsequently cloned using the restriction endo-nuclease sites *Bst*WI and *Bst*BI.

EXAMPLE 2. EXPRESSION OF HUMAN INTERFERON β IN A BACULOVIRUS/INSECT CELL SYSTEM

15 In order to express the synthetic gene, encoding human interferon β h harboured in pCBProF1 (described in example 1) in the baculovirus/insect cell system the gene was excised with *Nhe*I and *Xho*I and ligated into the transfer vector pBlueBac 4.5, which is included in the MaxBac 2.0 Transfection kit obtained from Invitrogen (San Diego, USA). All methods used for generation of recombinant baculovirus and expression in insect cells are described in the
20 "MaxBac 2.0 Transfection and Expression Manual" included in the kit.

In brief, together with linarized AcMNPV DNA (Bac-N-Blue DNA) pBlueBac 4.5-interferon β CBProF1) was transfected into SF9 cells. 3 days post-transfection the transfection supernatant was harvested and a plaque assay with appropriate viral dilutions was prepared. Blue distinct plaques were visible after 7 days and 6 individual plaques were collected
25 for propagation in a 6-well plate. After 5 days 2 ml virus supernatant (P-1 stock) was harvested from each well. 0.75 ml was taken out from the P1 stocks and viral genomic DNA was isolated. The viral genomic DNA's were analysed in PCR reactions with forward/reverse primers in order to be able to select the recombinant baculoviruses among the six P-1 stocks. A small aliquot from the recombinant P-1 stock was tested in a human interferon β specific ELISA (available
30 from PBL Biomedical Laboratories) in order to ensure that recombinant human interferon β was present in the supernatant.

For further propagation of chosen recombinant baculovirus 6×10^6 SF9 cells were seeded in a T-80 culture flask and infected with 200 μ l of the P-1 stock. After 5 days the supernatant (P-2 stock) was harvested and 2 ml of the P-2 stock was used to infect a 100 ml

5 suspension culture (1×10^6 SF9 cells/ml) in a 500 ml Erlenmeyer flask (Corning). After 5 days the supernatant (P-3 stock) was harvested and the virus titer was determined by plaque assay.

In order to produce human interferon β for purification 1×10^9 SF9 cells were harvested from a backup suspension culture. In a 50 ml screw-cap tube the SF9 cells were infected with recombinant baculovirus from the P-3 stock (MOI = 2) in a period of 15 minutes.

10 Hereafter the cells were spun down and washed one time in serum-free medium (Sf-900 II SFM, Gibco BRL) and transferred to a 2800 ml Triple Baffle Fernbach Flask (Bellco) containing 1 l serum-free medium. 3 days post-infection the medium supernatant was harvested and the recombinant human interferon β was purified.

Purification of interferon β molecules

15 The fermentation broth is concentrated and/or pH adjusted to approximately 4.5 after dilution to suitable ionic strength. Suitable is intended to mean that the ionic strength is so low that interferon β will bind to a Mono S cation exchange column (Pharmacia) equilibrated in 4 mM acetic acid pH 4.5 (buffer A). After application, the column is washed with 3 column volumes buffer A and interferon β is eluted with a linear gradient from buffer A to buffer A including 1 M NaCl. Alternatively purification can be obtained as described for Interferon α (Analytical Biochemistry 247, 434-440 (1997) using a TSK-gel SP-5PW column (Toso Haas)).

20 Alternatively His tagged interferon β can be purified using IMAC (Immobilized Metal Affinity Chromatography) in accordance with well known methods, e.g., as described by UniZyme Laboratories, Denmark.

25 Another purification method makes use of monoclonal or polyclonal antibodies. Interferon β fermentation broth is adjusted to pH 7 and 0.5 M NaCl and applied to a column with immobilized monoclonal antibody to recombinant human interferon β . The column is equilibrated with e.g. 10 mM Tris, 0.5 M NaCl, pH 7 (Buffer B) prior to application. After application the column is washed with 3 column volumes Buffer B and eluted with a suitable
30 buffer at low pH (e.g. pH 2-3).

Alternatively, if the interferon β is tagged with e.g. the c-Myc peptide (EQKLI SEEDL), monoclonal antibodies raised against the c-Myc peptide, can be used in a similar fashion. Immobilization of antibody to the column is achieved using e.g. CNBr-Sepharose (Pharmacia) according to the manufacturers instructions.

5 A combination of Cation exchange chromatography, IMAC and/or antibody chromatography may be applied if necessary to obtain relevant purity for further experiments.

Purity, identity, quantity and activity of eluted fractions from the abovementioned columns can be determined using a combination of methods known by the person skilled in the art. These may include one or more of the following assays and methods or other relevant
10 methods known by the person skilled in the art: the primary and secondary assays described above, ELISA methods, SDS-PAGE, western blotting, IEF, HPLC, amino acid sequencing, mass spectrometry and amino acid analysis.

Following purification, the modified interferon β polypeptide may be subjected to conjugation to a polymer molecule such as M-SPA-5000 from Shearwater Polymers according to
15 the manufacturer's instructions. Preferably, the receptor recognition site of the purified modified interferon β polypeptide is blocked prior to conjugation as described in the Materials and Methods section herein.

EXAMPLE 3. EXPRESSION OF HUMAN INTERFERON β IN HEK293 CELLS

In order to express the synthetic gene, encoding human interferon β , harboured by
20 pCBProF1 (described in example 1), in HEK293 cells (ATCC Cat. No. CRL-1573) the gene was PCR-amplified with the two primers PBR 7 (5'-CGCGGATCCATATGACCAACAAGTGCCTG-3') (SEQ ID NO 28) and PBR 2 (5'-CGCGGATCCTTATCAGTTGCGCAG-3') (SEQ ID NO 29) and cloned into the BamHI site of pcDNA3.1 (-) (Invitrogen, USA) in correct orientation, giving the plasmid pPR9.

25 For transfection of the HEK293 cell line a T-25 culture flask was seeded to 50% confluency in DMEM medium (Life Technologies, USA) containing 10% FBS and incubated over night. By usage of FuGENE 6 Transfection Reagent (Roche, USA) pPR9 was transfected into the cells: To 95 μ l serum-free DMEM medium was added 5 μ l FuGENE 6 and 1.7 μ l (2 μ g) pPR9 and incubated at room temperature for 20 minutes. The transfection complex was then
30 added drop-wise to the cells and the culture flask was returned to the incubator. Next day the cells were trypsinized and seeded into a T-80 culture flask in DMEM medium containing 10% FBS and 500 μ g Geneticin (Life Technologies) per ml.

At confluency it was confirmed, by usage of a human interferon β specific ELISA, that the primary transfection-pool was expressing the wished protein and the cells were

5 sub-cloned by limited dilution. In this way a high-producing HEK293 clone was identified expressing human interferon β .

EXAMPLE 4. HIGH LEVEL EXPRESSION OF INTERFERON β IN CHO CELLS

10 The cell line CHO K1 [p22]-E4 (ATCC # CCL-61) stably expressing human interferon β was passed 1:10 from a confluent culture and propagated as adherent cells in T-25 flasks in serum containing medium (MEM α w/ ribonucleotides and deoxyribonucleotides (Gibco/BRL Cat # 32571), 10% FCS (Gibco/BRL Cat # 10091), penicillin and streptomycin (Gibco/BRL Cat # 15140-114) until confluence. The media was then changed to serum free media (RenCyte CHO; MediCult Cat.# 22600140) for 24 hours before including 5 mM Sodium Butyrate (Merck Cat # 8.17500) during a medium change. The cells were then allowed to
15 express interferon β for 48 hours prior to harvest of the medium. The interferon β concentration in the duplicate cultures were determined to be 854,797 IU/ml (with lower and upper 95% confidence interval at 711,134 IU/ml and 1,032,012 IU/ml) respectively).

In a separate set of experiments, the cell line CHO K1 [p22]-E4 stably expressing human interferon β was passed 1:10 from a confluent culture and propagated as adherent cells in
20 serum containing medium (MEM α w/ ribonucleotides and deoxyribonucleotides (Gibco/BRL Cat # 32571), 10% FCS (Gibco/BRL Cat # 10091), penicillin and streptomycin (Gibco/BRL Cat # 15140-114) until confluence in a 10 layer cell factory (NUNC #165250). The media was then changed to serum free media; DMEM/F12 (Gibco/BRL # 11039-021) with the addition of 1:100 ITS-A (Gibco/BRL # 51300-044) and 1:500 EX-CYTE VLE (Serological Proteins Inc. # 81-
25 129-1) and 1:100 penicillin and streptomycin (Gibco/BRL Cat # 15140-114) for 48 hours before changing the medium with the further addition of 5 mM butyrate (Merck Cat # 8.17500). The cells were then allowed to express interferon β for 48 hours prior to harvest of the medium. The interferon β concentration was determined to be 824,791 IU/ml (with lower and upper 95% confidence interval at 610,956 IU/ml and 1,099,722 IU/ml) respectively).

30 It is contemplated that interferon β polypeptides of the invention may be produced in equally high yields in the same manner as any of those described above.

5 EXAMPLE 5. CONSTRUCTION AND EXPRESSION OF INTERFERON β VARIANT WITH ONE INTRODUCED GLYCOSYLATION SITE

In order to insert an extra N-linked glycosylation site at position 111 in hINF- β , the synthetic gene (*hinf- β*) encoding hINF- β (described in example 1) was altered by site-directed PCR mutagenesis. Using BIO-X-ACT (Bioline, UK) and the plasmid PF050 [*hinf- β* /pcDNA3.1(-)Hygro/Intron (a derivative of pcDNA3.1(-)Hygro (Invitrogen, USA) in which a chimeric intron obtained from pCI-neo (Promega, USA) had been inserted between the BamHI and NheI sites in the MCS of the vector] as template, two PCR reactions were performed with two overlapping primer-sets [CB41 (5'-TTTAAACTGGATCCAGCCACCATGACCAACAAG-3') (SEQ ID NO 30) /CB55 (5'-CGGCCATAGT

10 AGCGCTTCAGGTGCAGGGAGCTCATCAGCTTGCCGGTGGTGTGTGCTCCTCCTC-3') (SEQ ID NO 31) and CB42 (SEQ ID 26) / CB86 (5'-GAAGGAGGACAACACCACCGCAAGCTGATGAGCTCCCTGCACCTGAAGCGCTACT

15 ATGGCC G-3') (SEQ ID NO 32) resulting in two fragments of 446 and 184 base pairs, respectively. These two fragments were assembled in a third PCR with the flanking primers CB41 and CB42. The resulting gene was inserted into the mammalian expression vector pcDNA3.1(-)Hygro/Intron and confirmed by DNA sequencing to have the correct base changes leading to the substitutions F111N and R113T in hINF- β (plasmid designated PF085).

20

To test the activity of the [F111N+ R113T]hINF- β variant, PF085 was transfected into the CHO K1 cell line (ATCC #CCL-61) by use of Lipofectamine 2000 (Life Technologies, USA) as transfection agent. 24 hours later the culture medium was harvested and assayed for INF- β activity/concentration:

25

Activity: 56046 IU/ml [primary assay]
ELISA: 80 ng/ml
Specific activity: 7×10^8 IU/mg

30 As seen, the [F111N+R113T]hINF- β variant has a very high specific activity, about twice the specific activity of wt hINF- β .

5 EXAMPLE 6. CONSTRUCTION AND EXPRESSION OF INTERFERON β WITH
ANOTHER INTRODUCED GLYCOSYLATION SITE [Q49N+Q51T]

Analogously to what is described in Example 5 an extra N-linked glycosylation site was introduced in position 49 by means of the substitutions Q49N and Q51T. Using PF043 (*hinf*- β /pcDNA3.1 (Invitrogen, USA)) as template, two PCR reactions were performed with two overlapping primer-sets [PBR7 (SEQ ID NO 28) /PBR78 (5'-GGCGTCCTCCTTGGTGAAGTTCTGCAGCTG-3') (SEQ ID NO 33) and PBR8 (5'-ATATATCCCAAGCTTTTATCAGTTGCGCAGGTAGCCGGT-3') (SEQ ID NO 34)/PBR77 (5'-CAGCTGCAGAACTTCACCAAGGAGGACGCC-3') (SEQ ID NO 35) resulting in two fragments of 228 and 369 base pairs, respectively. These two fragments were assembled in a third PCR with the flanking primers PBR7 and PBR8. The resulting gene was inserted into the mammalian expression vector pcDNA3.1(-)Hygro/Intron and confirmed by DNA sequencing to have the correct base changes leading to [Q49N,Q51T]hINF- β (plasmid designated PF104).

To test the activity of the [Q49N+Q51T]hINF- β variant, PF104 was transfected into the CHO K1 cell line by use of Lipofectamine 2000 (Life Technologies, USA) as transfection agent. 24 hours later the culture medium was harvested and assayed for INF- β activity/concentration:

Activity:	17639 IU/ml [primary assay]
ELISA:	10 ng/ml
Specific activity:	1.7×10^9 IU/mg

As observed here the [Q49N+Q51T]hINF- β variant has a high specific activity. This may be due to poor recognition by one of the monoclonal antibodies used in the ELISA.

EXAMPLE 7. CONSTRUCTION AND EXPRESSION OF INTERFERON β WITH TWO INTRODUCED GLYCOSYLATION SITES

The additional glycosylation sites described in Examples 5 and 6 were introduced into human interferon β by means of the substitutions Q49N, Q51T, F111N, and R113T.

Using PF085 (described in example 5) as template, two PCR reactions were performed with two overlapping primer-sets [PBR89 (5'-CGCGGATCCAGCCACCATGACCAACAAGTGCCTG) (SEQ ID NO 36)/PBR78 (SEQ ID NO 33) and PBR8 (SEQ ID NO 34) /PBR77 (SEQ ID NO 35)] resulting in two fragments of 228 and 369 base pairs, respectively.

5 These two fragments were assembled in a third PCR with the flanking primers PBR89 and PBR8. The resulting gene was inserted into the mammalian expression vector pcDNA3.1(-)Hygro/Intron and confirmed by sequencing to have the correct base changes leading to [Q49N, Q51T, F111N, R113T] hINF- β (plasmid designated PF123).

10 PF123 was transfected into CHO K1 cells by use of Fugene 6 (Roche) as transfection agent. 24 hours later the culture medium was harvested and assayed for INF- β activity/concentration:

Activity:	29401 IU/ml	[primary assay]
ELISA:	14 ng/ml	
Specific activity:	2.1×10^9 IU/ml	

15 As observed here the [Q49N+Q51T+ F111N+ R113T]hINF- β variant also has a high specific activity.

 The variant was found to have receptor binding activity in the receptor binding assay described in the Materials and Methods section, which is based on the use of the crosslinking agent DSS.

20 EXAMPLE 8. PRODUCTION OF [Q49N+ Q51T+ F111N+ R113T]INTERFERON- β GLYCOSYLATION VARIANT IN ROLLER BOTTLES

 A CHOK1 sub-clone (5/G-10) producing the [Q49N+Q51T+F111N+R113T] glycosylation variant was seeded into 2 roller bottles, each with an expanded surface of 1700 cm² (Corning, USA), in 200 ml DMEM/F-12 medium (LifeTechnologies; Cat. # 31330) supplemented with 10% FBS and penicillin/streptomycin (P/S). After 2 days the medium was exchanged. After another 2 days the two roller bottles were nearly 100% confluent and the medium was shifted to 300 ml serum-free UltraCHO medium (BioWhittaker; Cat. # 12-724) supplemented with 1/500 EX-CYTE (Serologicals Proteins; Cat. # 81129N) and P/S. Growing the cells in this medium promotes a higher cell mass, higher than can be achieved in the serum containing medium. After 2 days the medium was renewed. After another 2 days the medium was shifted to the production medium: DMEM/F-12 medium (Life Technologies; Cat. # 21041) supplemented with 1/100 ITSA (Life Technologies; Cat. # 51300-044) [ITSA stands for Insulin (1.0 g/L) – Transferrin (0.55 g/L) – Selenium (0.67 mg/L) supplement for Adherent cultures], 1/500 EC-CYTE and P/S. In the figure below is shown the production run where 300 ml medium

5 was harvested from each roller bottle every day. The harvested media from the two roller bottles were pooled before a medium sample was taken out for interferon β activity determination. As seen in fig. 2 the production run was terminated after 26 days. After a lag-period of 5 days the activity mediated by the [49N+ Q51T+F111N+R113T]Interferon- β variant increased dramatically and for the rest of the production run the harvested interferon- β activity per day, in
10 average, was 2.4 million IU/ml x 600 ml = 1.440 billion IU. In total 3.2×10^{10} IU was produced corresponding to 160 mg protein (with a hypothetical specific activity of 2×10^8 IU/mg).

EXAMPLE 9. PRODUCTION, PURIFICATION, AND PEGYLATION OF THE INTERFERON- β VARIANT K19R+ K45R+ K123R

To end up with 100 ml serum-free medium containing the Interferon- β variant
15 K19R+K45R+K123R, 3 T-175 flasks were seeded with COS-7 cells in DMEM medium (Life technologies; Cat. # 21969-035) supplemented with 10% FBS plus Glutamine and penicillin/streptomycin. On the day of transfection (at nearly 100% confluency) the medium was renewed with 30 ml fresh medium 4 – 5 hours before the transfection. To prepare the transfection, 1890 μ l DMEM medium without supplements was aliquoted into a 14 ml
20 polypropylene tube (Corning). 210 μ l Fugene 6 (Roche) was added directly into the medium and incubated for 5 min at RT. In the meantime 168 μ g plasmid DNA ([K19R, K45R, K123R]INF- β /pcDNA3.1(-)Hygro; PF #161) was aliquoted into another 14 ml polypropylene tube. After 5 min incubation the Fugene 6 mix was added directly to the DNA solution and incubated for 15 min at RT. After incubation about 700 μ l was added drop wise to each of the three cell media.

25 Next day the transfection medium was substituted with 35 ml serum-free production medium. The serum-free medium is based on DMEM medium (Life Technologies; Cat. # 31053-028) supplemented with Glutamine, Sodium Pyruvate, penicillin/streptomycin, 1% ITSA (Life Technologies; Cat. # 51300-044), and 0.2% Ex-Cyte (Serologicals Proteins; Cat. # 81-129). Before the production medium was added the cell layers were washed two times in the
30 DMEM medium without additives.

Three days post-transfection the 100 ml serum-free medium was harvested for purification and PEGylation of the Interferon- β variant. pH was adjusted to 6.8 and conductivity adjusted to < 10 mS/cm with Milli Q water. Then the broth was batch adsorbed to 1 ml SP 550 cation exchange resin (TosoHaas) preequilibrated with buffer A (20 mM phosphate, 100 mM

NaCl, pH 7). After 2 h rotation end over end, the resin was allowed to sediment and transferred to a column. The resin was washed with 5 column volumes buffer A and eluted with 2 ml buffer B (20 mM phosphate, 800 mM NaCl, pH 7). The eluate was concentrated to 500 ul on VivaSpin (cutoff 10 kDa) after addition of 5 % ethyleneglycol. The concentrate was adjusted to 50 mM phosphate, 0.3 M NaCl, 20 % ethyleneglycol, pH 8 in a final volume of 2 ml and further concentrated to 0.5 ml.

The final concentrate was PEGylated as follows: to 100 ul of the final concentrate, 25 ul of activated mPEG-SPA (5000 kDa, Shearwater, Alabama) freshly prepared in phosphate buffer, pH 8 were added to make final concentrations of activated PEG of 0, 5, 10, 25 or 50 mg/ml. The reaction was allowed to proceed for 30 min at room temperature and then quenched by addition of 50 mM glycine buffer. Samples were frozen immediately at -80 C and bioactivity was measured as described (Primary Assay). Western blots of each sample were performed in order to evaluate the amount of unreacted Interferon- β variant present in the PEGylated sample.

Results demonstrate that at 25 mg activated PEG/ml, nonPEGylated Interferon- β variant was absent as judged by western blot and the variant retained 50 % of its bioactivity compared to the control sample (treated identically, but with 0 mg/ml activated PEG).

EXAMPLE 10. EXPRESSION AND PURIFICATION OF SOLUBLE IFNAR2

The cDNA's encoding the extracellular domain of IFNAR-1 and IFNAR-2 (termed IFNAR1ec and IFNAR2ec, respectively) were amplified from HeLa cell cDNA using PCR with primers corresponding to the first 10 amino acid residues and the final 10 amino acid residues of the extracellular domain of IFNAR-2 (the nucleotide sequence of which is apparent from Novick et al., Cell, Vol. 77, pp 391-400, 1994) and the first 10 amino acid residues and the final 10 amino acid residues of the extracellular domain of IFNAR-1 (the nucleotide sequence of which is apparent from Uze et al., Cell Vol. 60, 225-234, 1990). The cDNA's were subcloned into the pBlueBac 4.5/V5-His-TOPO vector (Invitrogen) and a recombinant Baculovirus obtained by homologous recombination, plaque purification, and propagation in Sf9 cells. Sf9 cells were infected with the recombinant Baculovirus and expression from the resulting cells was obtained essentially as described in Example 2.

IFNAR1ec and IFNAR2ec protein was observed in culture supernatants two to three days after infection of Sf9 cells with recombinant baculovirus. The activity of soluble receptors was observed in an Interferon antagonist assay. Briefly, Hela cells containing the ISRE element (as described in the Primary Assay above) are stimulated with a sub-maximal dose of human wild-type Interferon β in the presence of varying concentrations of IFNARec supernatant. The antagonist effect of the supernatant is directly proportional to the amount of soluble receptor present.

IFNAR2ec was purified from filtered culture supernatants using ion exchange, and affinity chromatography. Culture supernatants positive for IFNAR2ec were pH adjusted to 7.5 and loaded onto an anion exchange column, and the bound recombinant protein was eluted using 500mM NaCl. The partially pure IFNAR2ec was then diluted and pH adjusted to 8.0, before further purification using binding to a TALON™ Metal Affinity Resin and elution with imidazol. The final preparation was frozen in aliquots. IFNAR1ec can be purified as described for IFNAR2ec with the exception that cation exchange chromatography at pH6.0 will be used as the ion exchange step.

EXAMPLE 11. USE OF SOLUBLE IFNAR2 FOR PURIFICATION AND PEGYLATION OF INTERFERON- β AND VARIANTS THEREOF

Purified IFNAR2 obtained as described in Example 9 is immobilized either through amino or carboxyl groups using e.g. CNBr-activated Sepharose 4B or EAH Sepharose 4B according to the manufacturer's instructions (Amersham Pharmacia Biotech, Affinity Chromatography, Principles and Methods, 18-1022-29, edition AB). It is critically important that the coupling method allows functional IFNAR2 to be immobilized and this is tested through optimization of the coupling conditions (pH, coupling buffer, ratio of IFNAR 2 to activate matrix etc). Another critical parameter is the blocking of excess active groups. Subsequently, testing of binding capacity by addition of interferon- β and measurement of breakthrough is carried out.

Optimally immobilized IFNAR2 is used for purification of Interferon- β as follows. A 5 ml column with 1 mg IFNAR 2 immobilized per ml gel is equilibrated with buffer A (20 mM phosphate, 300 mM NaCl, pH 7). Then the column is loaded with a 2 mg Interferon- β sample in buffer A and subsequently washed with 5 column volumes buffer A. Elution is obtained by pumping 2 column volumes of buffer B onto the column. Fractions of 1 ml are collected and assayed for bioactivity. Optimal elution conditions are dependent on the

5 immobilization method, but examples of elution conditions include pH 1.5 – 3 (e.g. 0.1 M glycine pH 2.3 in 0.5 M NaCl), pH 11.5 – 12, 3.5 M MgCl₂, 6M urea or the like.

EXAMPLE 12. USE OF IMMOBILIZED IFNAR2 FOR PEGYLATION OF INTERFERON β (VARIANTS)

10 In addition to the use described in Example 10, immobilized IFNAR 2 may be used for optimal PEGylation, wherein PEGylation of the part of Interferon- β or variants thereof interacting with the receptor is avoided.

A 5 ml column with 1 mg IFNAR 2 immobilized per ml gel is equilibrated with buffer A (20 mM phosphate, 300 mM NaCl, pH 7). Then the column is loaded with a 2 mg Interferon- β sample in buffer A and subsequently washed with 5 column volumes buffer A. A
15 solution of activated mPEG-SPA (1–50 mg/ml in buffer A) is pumped on the column and allowed to react for 15 min–12 h depending on temperature. One preferred range of combination of residence time and temperature is 15–60 min, 10–20 °C, another is 30 min to 5 h, 2–8°C . After the indicated time period, elution is obtained by pumping 2 column volumes of buffer B onto the column. Fractions of 1 ml are collected and assayed for bioactivity using the primary
20 screening assay. Optimal elution conditions are dependent on the immobilization method, but examples of elution conditions include pH 1.5–3 (e.g. 0.1 M glycine pH 2.3 in 0.5 M NaCl), pH 11.5–12, 3.5 M MgCl₂, 6M Urea or the like.

EXAMPLE 13. ANTIVIRAL ACTIVITY OF PEGYLATED VARIANT

25 The pegylated IFN- β variant protein, K19R+K45R+K123R, was assayed using the antiviral bioassay. Wild-type and variant proteins were added to A549 cells in concentrations from 10⁻¹⁰–0.0001 IU/mL in triplicate cultures.

The pegylated IFN- β variant showed total inhibition of EMC virus induced cell death at a concentration of 3 IU/mL, with an EC₅₀ of 0.13 IU/mL (Figure 1). The wild-type standard shows virus inhibition with an EC₅₀ of 1.4 IU/mL.

30 These results demonstrate that the pegylation of the modified interferon β polypeptide resulted in a conjugate with full anti-viral activity.

5 EXAMPLE 14. ANTIBODY NEUTRALISATION OF GLYCOSYLATED VARIANT

The antibody neutralisation of wild-type and glycosylated IFN- β variant protein, Q49N+Q51T+F111N+R113T, was assayed using the ISRE neutralisation assay. Interferon β wild-type and variant proteins (in five fold dilutions starting at 12500 IU/mL) were incubated with polyclonal rabbit anti-interferon β antibody (PBL Biomedical Laboratories) in
 10 concentrations 0, 40 and 200 ng/mL.

In the presence of 200 ng/mL polyclonal rabbit anti-serum the activity of the wild type interferon β protein was reduced 11.8 times whereas the activity of the glycosylated interferon β variant only was reduced 3.0 times. Thus the degree of antibody recognition of the interferon β variant was reduced by 75% of the wt level, see Table 1 below. These results
 15 demonstrate that the recognition of the glycosylated mutant interferon β by polyclonal antibodies raised in animals immunised with wild-type human interferon β is highly reduced. Thus, a large portion of the immunogenic epitopes in wild-type human interferon β have been removed/shielded by the modifications made in the variant molecule.

Table 1

Antibody conc. (ng/mL)	Protein	EC50	Fold inhibition	Reduction of antibody neutralisation
0	wt	0.00039	-	-
	variant	0.00020	-	-
40	wt	0.00190	4.8	-
	variant	0.00020	1.0	79%
200	wt	0.00461	11.8	-
	variant	0.00059	3.0	75%

20 EXAMPLE 15. CONSTRUCTION AND EXPRESSION OF INTERFERON β MOLECULES WITH MODIFIED N-TERMINAL

N-terminally modified variants of interferon β were constructed as described in the preceding examples.

25 For the construction of an expression plasmid for the interferon β variant, INFB S(-1)A + M1Q the following primers were used:

CBProFpr110:

AAC TGG ATC CAG CCA CCA TGA CCA ACA AGT GCC TGC TCC AGA TCG CCC TGC TCC TGT GCT TCA GCA CCA CGG CCC TAG CCC AGA GCT AC (SEQ ID NO 37) and

30 CBProFpr42 (SEQ ID NO 26),

5 For the construction of an expression plasmid for the interferon variant, INF β S(-1)AQ (indicating substitution of the S residue located in position (-1) with an A and a Q residue) the following primers were used:

CBProFpr109:

AAC TGG ATC CAG CCA CCA TGA CCA ACA AGT GCC TGC TCC AGA TCG CCC TGC

10 TCC TGT GCT TCA GCA CCA CGG CCC TAG CCC AGA TGA GCT AC (SEQ ID NO 38)

and

CBProFpr42 (SEQ ID NO 26).

To test the activity of these variants the respective plasmids; pF154 and pF163 were transfected into CHO K1 cells using Lipofectamine 2000 (Life Technologies, USA) as
15 transfection reagent. The supernatants were harvested 24 hours post transfection and assayed in the primary activity assay and in the ELISA as described in the Materials and Methods section.

The following results were obtained:

INFB S-1A + M1Q (pF154):

Activity: 106410 IU/ml

20 ELISA: 333 ng/ml

Specific activity: 3.2×10^8 IU/mg

INFB S-1AQ (pF163):

Activity: 90634 IU/ml

25 ELISA: 193 ng/ml

Specific activity: 4.7×10^8 IU/mg

These molecules are as active as wild type human interferon β .

EXAMPLE 16. PREPARATION OF PEGYLATED IFN- β VARIANTS

50 microliters of a 0.3 mg/ml solution of recombinant human IFN- β polypeptide comprising the mutations Q49N+Q51T+ K19R+ K45R+ K123R in 50 mM Na-acetate, 35%
30 ethylene glycol, pH 5.5 were mixed with 10 μ l 0.5 M Na-phosphate, pH 8.0 and 20 μ l 50 mM Na-phosphate, 0.1 M NaCl, 30% ethylene glycol, pH 8.0 containing 0.02 mg/ml SPA-mPEG (N-succinimidyl Propionate methoxy polyethylene glycol). This corresponds to a 10 molar excess of SPA-mPEG to IFN- β .

After $\frac{1}{2}$ hour with gentle rotation at room temperature, the reaction was quenched
35 by addition of 5 μ l 20 mM Glycine, pH 8.0. At this stage, the reaction mixture contained a mixture of unmodified as well as pegylated forms of recombinant human IFN- β .

005220-159584960

5

Activity:

In vitro testing using the primary screening assay demonstrated that the pegylated material retained 40 % activity, as compared to the unmodified recombinant human IFN- β .

10 In another experiment, 50 μ l of a 0.14 mg/ml solution of recombinant human IFN β polypeptide comprising the mutations Q49N+Q51T in 50 mM Na-acetate, 35% ethylene glycol, pH 5.5 was mixed with 10 μ l 0.5 M Na-phosphate, pH 8.0 and 20 μ l 50 mM Na-phosphate, 0.1 M NaCl, 30% ethylene glycol, pH 8.0 containing 0.03 mg/ml SPA-mPEG. This gave a 10 molar excess of SPA-mPEG to IFN- β .

15 After $\frac{1}{2}$ hour with gentle rotation at room temperature, the reaction was quenched by addition of 5 μ l 20 mM Glycine, pH 8.0. At this stage, the reaction mixture contained a mixture of unmodified as well as pegylated forms of recombinant human IFN- β .

Activity:

In vitro testing using the primary screening assay demonstrated that the pegylated material retained 20 % activity, as compared to the unmodified recombinant human IFN- β .

20

25

While the foregoing invention has been described in some detail for purposes of clarity and understanding, it will be clear to one skilled in the art from a reading of this disclosure that various changes in form and detail can be made without departing from the true scope of the invention. For example, all the techniques, methods, compositions, apparatus and systems described above may be used in various combinations. *All publications, patents, patent applications, or other documents cited in this application are incorporated by reference in their entirety for all purposes to the same extent as if each individual publication, patent, patent application, or other document were individually indicated to be incorporated by reference for all purposes.*

5 WHAT IS CLAIMED IS:

1. A conjugate exhibiting interferon β activity, which conjugate comprises at least one first non-polypeptide moiety covalently attached to an interferon β polypeptide variant, which interferon β polypeptide variant differs from a wild-type human interferon β by at least one introduction and at least one removal of an amino acid residue, which amino acid residue
10 comprises an attachment group for the first non-polypeptide moiety.

2. The conjugate of claim 1, wherein the first non-polypeptide moiety is selected from the group consisting of: a polymer molecule, a lipophilic compound, a sugar moiety and an organic derivatizing agent.

3. The conjugate of claim 2, wherein the polymer molecule comprises linear or
15 branched polyethylene glycol.

4. The conjugate of claim 1, wherein the first non-polypeptide moiety comprises a polymer molecule having an attachment group selected from the group consisting of: lysine, aspartic acid, glutamic acid and cysteine.

5. The conjugate of claim 4, wherein the first non-polypeptide moiety is a
20 polymer molecule having lysine as an attachment group.

6. A conjugate exhibiting interferon β activity, which conjugate comprises at least one first non-polypeptide moiety conjugated to at least one lysine residue of an interferon β polypeptide variant, which interferon β polypeptide variant differs from a wild-type human interferon β by at least one introduction or removal of a lysine residue.

7. The conjugate of claim 6, which interferon β polypeptide variant differs
25 from a wild-type human interferon β by introduction and removal of a lysine residue.

8. The conjugate of claim 6, wherein the lysine residue removed is selected from one or more residue selected from the group consisting of: K19, K33, K45, K52, K99, K105, K108, K115, K123, K134 and K136.

5 12. The conjugate of claim 11, wherein the interferon β polypeptide variant comprises at least one substitution selected from the group consisting of: N4K, R11K, G26K, R27K, Q48K, Q49K, R71K, D73K, S75K, E85K, A89K, Y92K, H93K, F111K, R113K, L116K, R124K, G127K and Y155K.

10 13. The conjugate of claim 12, wherein the substitution is selected from the group consisting of: Q49K and F111K.

14. The conjugate of claim 6, comprising at least two introduced lysine residues.

15 15. The conjugate of claim 11, wherein the interferon β polypeptide variant further comprises the removal of at least one lysine residue.

16. The conjugate of claim 15, wherein the at least one lysine residue is selected from the group consisting of: K19, K33, K45, K52, K99, K105, K108, K115, K123, K134 and K136.

17. The conjugate of claim 15, comprising one of the following sets of mutations:

20 K19R+K45R+F111K+K123R;
K19R+K45R+Q49K+F111K+K123R;
K19R+K45R+Q49K+K123R;
K19R+K45R+ F111K;
K19R+K45R+Q49K+F111K;
25 K19R+Q49K+K123R;
K19R+Q49K+F111K+K123R;
K45Q+F111K+K123Q;
K45R+Q49K+K123R; or
K45R+Q49K+F111K+K123R.

30 18. The conjugate of claim 1 or 6, wherein the non-polypeptide moiety comprises a polymer selected from the group consisting of: SS-PEG, NPC-PEG, aldehyd-PEG, mPEG-SPA, PEG-SCM and mPEG-BTC.

19. A conjugate exhibiting interferon β activity, which conjugate comprises at least one first non-polypeptide moiety conjugated to at least one cysteine residue of an interferon β polypeptide variant, which interferon β polypeptide variant differs from a wild type interferon

5 β polypeptide by the introduction of at least one cysteine residue at a position selected from the group consisting of: F8, L9, R11, S12, F15, Q16, Q18, L20, W22, L28, L32, M36, P41, T58, Q64, N65, F67, I83, E85, N86, A89, N90, Y92, H93, H97, T100, L102, E103, L106, M117, L120, H121, R124, G127, R128, L130, H131, H140, I145, R147, V148, E149, R152, Y155, and F156 of SEQ ID NO 2.

10 20. The conjugate of claim 19, wherein the interferon β polypeptide variant further comprises removal of at least one cysteine residue.

21. The conjugate of claim 20, wherein the cysteine residue removed comprises C17.

15 22. The conjugate of claim 19, wherein the interferon β polypeptide variant comprises at least one of the mutations C17S or N80C.

23. The conjugate of claim 19, wherein the interferon β polypeptide variant comprises the mutations C17S and N80C.

24. The conjugate of claim 19, wherein the first non-polypeptide moiety is a polymer molecule.

20 25. A conjugate exhibiting interferon β activity, which conjugate comprises at least one first non-polypeptide moiety having an acid group as an attachment group, which moiety is conjugated to at least one aspartic acid or glutamic acid residue of an interferon β polypeptide variant, which interferon β polypeptide variant differs from a wild-type human interferon β by at least one introduction or removal of an aspartic acid or a glutamic acid residue.

25 26. The conjugate of claim 25, wherein at least one amino acid residue selected from the group consisting of: N4, L5, L6, F8, L9, Q10, R11, S12, S13, F15, Q16, Q18, K19, L20, W22, Q23, L24, N25, G26, R27, Y30, M36, Q46, Q48, Q49, I66, F67, A68, I69, F70, R71, S75, T82, I83, L87, A89, N90, V91, Y92, H93, Q94, I95, N96, H97, K108, F111, L116, L120, K123, R124, Y126, G127, R128, L130, H131, Y132, K134, A135, H140, T144, R147, Y155, 30 F156, N158, R159, G162, Y163 and R165 of SEQ ID NO 2 is substituted with an aspartic acid residue or a glutamic acid residue.

5 27. The conjugate of claim 25, comprising introduction of at least two introduced aspartic acid or glutamic acid residues.

 28. The conjugate of claim 25, comprising at least two first non-polypeptide moieties.

10 29. The conjugate of claim 25, wherein the first non-polypeptide moiety is a polymer molecule.

 30. The conjugate of claim 29, wherein the second non-polypeptide moiety comprises a sugar moiety.

15 31. The conjugate of claim 34, wherein the amino acid sequence of the interferon β polypeptide variant further comprises at least one of an introduction or a removal of an *in vivo* glycosylation site.

 32. The conjugate of claim 31, wherein the first non-polypeptide moiety is a polymer molecule having lysine as an attachment group.

 33. The conjugate of claim 1, 6, 19 or 25, which conjugate comprises a second non-polypeptide moiety.

20 34. The conjugate of claim 33, wherein the polypeptide comprises at least one removed amino acid residue comprising an attachment group for the first non-polypeptide moiety, and at least one introduced amino acid residue comprising an attachment group for the second non-polypeptide moiety.

25 35. The conjugate of claim 33, wherein the amino acid sequence of the interferon β polypeptide variant comprises at least two removed amino acid residues comprising attachment groups for the first non-polypeptide moiety and at least one introduced amino acid residue comprising an attachment group for the second non-polypeptide moiety.

 36. The conjugate of claim 33, wherein the second non-polypeptide moiety comprises a sugar moiety.

5 **37.** A conjugate exhibiting interferon β activity, which conjugate comprises at least one polymer molecule and at least one sugar moiety covalently attached to an interferon β polypeptide variant, which interferon β polypeptide variant differs from a wild-type human interferon β by

 (a) at least one introduction or removal of an amino acid residue, which amino
10 acid residue comprises an attachment group for the polymer molecule, and

 (b) at least one introduction or removal of an amino acid residue, which amino acid residue comprises an attachment group for the sugar moiety,

 with the proviso that when the attachment group for the polymer molecule is a cysteine residue, and the sugar moiety is an N-linked sugar moiety, introduction of a cysteine
15 residue does not destroy an N-glycosylation site.

38. The conjugate of claim 37, wherein the polymer molecule has lysine as an attachment group.

39. The conjugate of claim 38, wherein the polypeptide variant comprises at least one removed amino acid residue comprising an attachment group for the first non-
20 polypeptide moiety and at least one introduced amino acid residue comprising an attachment group for the second non-polypeptide moiety.

40. The conjugate of claim 37, wherein the interferon β polypeptide variant comprises one of the following sets of mutations:

 K19R+K45R+Q49N+Q51T+F111N+R113T+K123R;
25 K19R+K45R+Q49N+Q51T+F111N+R113T; or
 K19R+K45R+Q49N+Q51T+ K123R.

41. The conjugate of claim 1, 6, 19, 25 or 37, wherein the interferon β polypeptide variant comprises a modified N-terminus, which modified N-terminus is unavailable for conjugation to a non-polypeptide moiety.

30 **42.** A conjugate exhibiting interferon β activity, which conjugate comprises an interferon β polypeptide variant, which interferon β polypeptide variant differs from a wild-type human interferon β by introduction of at least one glycosylation site, the conjugate further comprising at least one un-PEGylated sugar moiety attached to the introduced glycosylation site.

5 **43.** A conjugate exhibiting interferon β activity, which conjugate comprises an
interferon β polypeptide variant, which interferon β polypeptide variant differs from a wild-type
human interferon β by at least one introduction or removal of a glycosylation site by introducing
or removing one or more amino acid residues comprising a glycosylation site in a position that is
occupied by an amino acid residue having at least 25% of its side chain surface exposed in
10 wildtype human interferon β .

44. The conjugate of claim 42 or 43, wherein the interferon β polypeptide
variant comprises at least one mutation selected from the group consisting of: S2N+N4T,
L9N+R11T, R11N, S12N+N14T, F15N+C16S, Q16N+Q18T, K19N+L21T, Q23N+H25T,
G26N+L28T, R27N+E29T, L28N+Y30T, D39T, K45N+L47T, Q46N+Q48T, Q48N+F50T,
15 Q49N+Q51T, Q51N+E53T, R71N+D73T, Q72N, D73N, S75N, S76N+G78T, L88T, Y92T,
N93N+I95T, L98T, E103N+K105T, E104N+L106T, E107N+E109T, K108N+D110T, D110N,
F111N+R113T and L116N.

45. The conjugate of claim 44, wherein the interferon β polypeptide comprises
one of the following sets of substitutions: Q49N+Q51T; Q49N+Q51T+F111N+R113T; or
20 Q49N+Q51T+R71N+D73T+F111N+R113T.

46. The conjugate of claim 42 or 43, wherein the amino acid sequence further
differs by removal of an amino acid comprising a glycosylation site.

47. The conjugate of claim 46, wherein the glycosylation site comprises an N-
glycosylation site.

25 **48.** A conjugate exhibiting interferon β activity, which conjugate comprises a
sugar moiety covalently attached to an interferon β polypeptide variant, which interferon β
polypeptide variant differs from a wild-type human interferon β by removal of at least one
glycosylation site.

49. The conjugate of claim 43 or 48, wherein an N-glycosylation site is removed
30 by the mutation N80C.

5 **50.** The conjugate of claim 1, 6, 19, 25, 37 or 43, wherein the interferon β polypeptide further comprises at least one substitution in a position selected from among M1, C17, N80 or V101.

51. The conjugate of claim 50, wherein the interferon β polypeptide variant comprises at least one substitution selected from M1del, M1K or C17S.

10 **52.** A nucleotide sequence encoding an interferon β polypeptide variant, which interferon β polypeptide variant comprises component of a conjugate exhibiting interferon β activity.

53. A host cell comprising the nucleotide sequence of claim 52.

54. An expression vector comprising the nucleotide sequence of claim 52.

15 **55.** A host cell comprising the expression vector of claim 54.

56. The host cell of claim 53 or 55, which host cell comprises a CHO cell, a BHK cell, a HEK293 cell or an SF9 cell.

57. A method of reducing immunogenicity or increasing functional *in vivo* half-life or serum half-life of an interferon β polypeptide variant, the method comprising:

20 (a) introducing an amino acid residue comprising an attachment group for a first non-polypeptide moiety into a position exposed at the surface of an interferon β polypeptide that does not contain such group; or

 (b) removing an amino acid residue comprising an attachment group for the first non-polypeptide moiety; and

25 conjugating the interferon β polypeptide variant with the first non-polypeptide moiety.

58. The method of claim 57, wherein the non-polypeptide moiety is selected from the group consisting of a polymer molecule, a sugar moiety, a lipophilic group and an organic derivatizing agent.

5 **59.** A method for preparing a conjugate exhibiting interferon β activity, the method comprising:
 contacting an interferon β polypeptide variant with a first non-polypeptide moiety under conditions conducive for conjugation; and
 recovering the conjugate exhibiting interferon β activity.

10 **60.** A pharmaceutical composition comprising a conjugate exhibiting interferon β activity and a pharmaceutically acceptable diluent, carrier, excipient or adjuvant.

61. The pharmaceutical composition of claim 60, wherein the pharmaceutical composition is effective for the treatment of one or more diseases.

15 **62.** The pharmaceutical composition of claim 61, wherein the pharmaceutical composition is effective for the treatment of multiple sclerosis.

63. A method for treating a mammal with multiple sclerosis comprising administering an effective amount of the pharmaceutical composition of claim 60.

20 **64.** A method for treating a mammal having circulating antibodies against interferon β 1a and/or 1b, which method comprises administering an effective amount of the pharmaceutical composition of claim 60.

65. The method of claim 63 or 64, wherein the mammal is a human.

66. A cell culture comprising
 a) a host cell transformed with a nucleotide sequence encoding an interferon β polypeptide variant; and,
25 b) a culture medium.

67. The cell culture of claim 66, wherein the culture medium comprises an interferon β polypeptide variant produced by expressing the nucleotide sequence.

68. The cell culture of claim 67, wherein the concentration of the interferon β polypeptide variant is at least 800,000 IU/ml of medium.

5 **69.** The cell culture of claim 67, wherein the concentration of the interferon β polypeptide variant is in the range of 800,000-3,500,000 IU/ml medium.

70. A cell culture comprising a host cell of claim 53 or 55.

71. A method of producing an interferon β polypeptide variant, the method comprising:

10 (a) culturing a cell expressing an interferon β polypeptide variant in a culture medium, such that the concentration of the interferon β polypeptide variant in the medium is at least 800,000 IU/ml medium; and

 (b) recovering the interferon β polypeptide variant.

15 **72.** The method of claim 71, wherein the concentration of the interferon β polypeptide variant is between 800,000 and 3,000,000 IU/ml medium.

ABSTRACT OF THE DISCLOSURE

The present invention provides new interferon β conjugates, methods of preparing such conjugates and the use of such conjugates in therapy, in particular for the treatment of multiple sclerosis.

5

Figure 1





Figure 2



005280" 63534260

DECLARATION

As a below named inventor, I declare that:

My residence, post office address and citizenship are as stated below next to my name; I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: **NEW INTERFERON β -LIKE MOLECULES** the specification of which X is attached hereto or was filed on as Application No. , and was amended on (if applicable).

I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, Section 1.56. I claim foreign priority benefits under Title 35, United States Code, Section 119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Country	Application No.	Date of Filing	Priority Claimed Under 35 USC 119
DK	PA 1999 01197	August 27, 1999	Yes
DK	PA 1999 01691	November 26, 1999	Yes
DK	PA 2000 00194	February 7, 2000	Yes
DK	PA 2000 00363	March 7, 2000	Yes
DK	PA 2000 00642	April 14, 2000	Yes

I hereby claim the benefit under Title 35, United States Code § 119(e) of any United States provisional application(s) listed below:

Application No.	Filing Date
60/160,782	October 21, 1999
60/169,077	December 6, 1999
60/189,599	March 15, 2000
60/202,248	May 5, 2000

I claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, Section 1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

Application No.	Date of Filing	Status

Full Name of Inventor 1:	Full Name: Anders Hjelholt Pedersen		
Residence & Citizenship:	City: Lyngby	State/Foreign Country: Denmark	Country of Citizenship: DK
Post Office Address:	Post Office Address: 58, Nybrovaenget, DK-2800 Lyngby DENMARK		
Full Name of Inventor 2:	Full Name: Hans Thalsgaard Schambye		
Residence & Citizenship:	City: Frederiksberg	State/Foreign Country: Denmark	Country of Citizenship: DK
Post Office Address:	Post Office Address: 21, Pelargonievej DK-2000 Frederiksberg DENMARK		
Full Name of Inventor 3:	Full Name: Kim Vilbour Andersen		
Residence & Citizenship:	City: Copenhagen	State/Foreign Country: Denmark	Country of Citizenship: DK
Post Office Address:	Post Office Address: 31, Taasingegade DK-2100 Copenhagen DENMARK		
Full Name of Inventor 4:	Full Name: Claus Bornaes		
Residence & Citizenship:	City: Hellerup	State/Foreign Country: Denmark	Country of Citizenship: DK
Post Office Address:	Post Office Address: 6, Hoejgaards Alle DK-2900 Hellerup DENMARK		
Full Name of Inventor 5:	Full Name: Paul Baad Rasmussen		
Residence & Citizenship:	City: Copenhagen	State/Foreign Country: Denmark	Country of Citizenship: DK
Post Office Address:	Post Office Address: 4, Borthigsgade DK-2100 Copenhagen DENMARK		

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor 1 Anders, Hjelholt Pedersen Date	Signature of Inventor 2 Hans, Thalsgaard Schambye Date	Signature of Inventor 3 Kim Vilbour Andersen Date
Signature of Inventor 4 Claus Bornaes Date	Signature of Inventor 5 Paul Baad Rasmussen Date	

I hereby certify that this is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated below, addressed to: Assistant Commissioner for Patents Washington, Box Patent Application, D.C. 20231, on August 25, 2000

LAW OFFICES OF JONATHAN ALAN QUINE

By

Tracie Brooks

Tracie Brooks

Attorney Docket No. 02-107600US

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

Anders Hjelholt Pedersen et al.

Application No.: Not Yet Known

Filed: Herewith

For: **NEW INTERFERON β -LIKE MOLECULES**

Examiner: Unassigned

Art Unit: Unassigned

STATEMENT ACCOMPANYING
SEQUENCE LISTING

Assistant Commissioner for Patents
Box Sequence Listing
Washington, D.C. 20231

Sir:

The undersigned hereby states that the Sequence Listing submitted concurrently herewith does not include matter which goes beyond the content of the application as filed and that the information recorded on the diskette submitted concurrently herewith is identical to the written Sequence Listing.

Respectfully submitted,

Gwynedd Warren

Gwynedd Warren, Ph.D.
Reg. No. 45,200

Aug. 25, 2000

Date

LAW OFFICES OF JONATHAN ALAN QUINE
P.O. BOX 458
Alameda, CA 94501
Ph.: (510) 337-7871 / Fax (510) 337-7877

005280" 69597950

5 SEQUENCE LISTING

<110> Maxygen ApS

<120> New Interferon Beta-like Molecules

<130> 2wo2

<140>

<141>

<160> 38

<170> PatentIn Ver. 2.1

<210> 1

<211> 840

<212> DNA

<213> Homo sapiens

<400> 1

```
acattctaac tgcaaccttt cgaagccttt gctctggcac aacaggtagt aggcgacact 60
gttcgtgttg tcaacatgac caacaagtgt ctccctccaa ttgctctcct gttgtgcttc 120
tccactacag ctctttccat gagctacaac ttgcttggtt tcctacaaag aagcagcaat 180
tttcagtgtc agaagctcct gtggcaattg aatgggaggc ttgaatactg cctcaaggac 240
aggatgaact ttgacatccc tgaggagatt aagcagctgc agcagttcca gaaggaggac 300
gccgcattga ccatctatga gatgctccag aacatctttg ctattttcag acaagattca 360
tctagcactg gctggaatga gactattgtt gagaacctcc tggctaattg ctatcatcag 420
ataaaccatc tgaagacagt cctggaagaa aaactggaga aagaagattt caccagggga 480
aaactcatga gcagtctgca cctgaaaaga tattatggga ggattctgca ttacctgaag 540
gccaaggagt acagtcactg tgccctggacc atagtcagag tggaaatcct aaggaacttt 600
tacttcatta acagacttac aggttacctc cgaaactgaa gatctcctag cctgtgcctc 660
tgggactgga caattgcttc aagcattctt caaccagcag atgctgttta agtgactgat 720
ggctaattga ctgcatatga aaggacacta gaagattttg aaatttttat taaattatga 780
gttattttta tttattttaa ttttattttg gaaaataaat tttttttggt gcaaaagtca 840
```

<210> 2

<211> 166

<212> PRT

<213> Homo sapiens

<400> 2

```
Met Ser Tyr Asn Leu Leu Gly Phe Leu Gln Arg Ser Ser Asn Phe Gln
  1             5             10             15

Cys Gln Lys Leu Leu Trp Gln Leu Asn Gly Arg Leu Glu Tyr Cys Leu
      20             25             30

Lys Asp Arg Met Asn Phe Asp Ile Pro Glu Glu Ile Lys Gln Leu Gln
      35             40             45

Gln Phe Gln Lys Glu Asp Ala Ala Leu Thr Ile Tyr Glu Met Leu Gln
      50             55             60

Asn Ile Phe Ala Ile Phe Arg Gln Asp Ser Ser Ser Thr Gly Trp Asn
      65             70             75             80
```

5 Glu Thr Ile Val Glu Asn Leu Leu Ala Asn Val Tyr His Gln Ile Asn
85 90 95

10 His Leu Lys Thr Val Leu Glu Glu Lys Leu Glu Lys Glu Asp Phe Thr
100 105 110

Arg Gly Lys Leu Met Ser Ser Leu His Leu Lys Arg Tyr Tyr Gly Arg
115 120 125

15 Ile Leu His Tyr Leu Lys Ala Lys Glu Tyr Ser His Cys Ala Trp Thr
130 135 140

Ile Val Arg Val Glu Ile Leu Arg Asn Phe Tyr Phe Ile Asn Arg Leu
145 150 155 160

20 Thr Gly Tyr Leu Arg Asn
165

25 <210> 3
<211> 70
<212> DNA
<213> Artificial Sequence

30 <220>
<223> Description of Artificial Sequence: primer

35 <400> 3
ggctagcggtt taaacttaag cttcgccacc atgaccaaca agtgcttgct ccagatcgcc 60
ctgctcctgt 70

40 <210> 4
<211> 70
<212> DNA
<213> Artificial Sequence

45 <220>
<223> Description of Artificial Sequence: primer

50 <400> 4
acaacctgct cggcttcctg cagaggagtt cgaacttcca gtgccagaag ctctgtggc 60
agctgaacgg 70

55 <210> 5
<211> 70
<212> DNA
<213> Artificial Sequence

60 <220>
<223> Description of Artificial Sequence: primer

<400> 5
gaacttcgac atccccgagg aaatcaagca gctgcagcag ttccagaagg aggacgccgc 60

5 tctgaccatc 70

<210> 6
<211> 70
10 <212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer

15 <400> 6
ttccgccagg actccagctc caccggttgg aacgagacca tcgtggagaa cctgctggcc 60
aacgtgtacc 70

20 <210> 7
<211> 70
<212> DNA
<213> Artificial Sequence

25 <220>
<223> Description of Artificial Sequence: primer

30 <400> 7
aggagaagct ggagaaggag gacttcaccc gcggcaagct gatgagctcc ctgcacctga 60
agcgtacta 70

35 <210> 8
<211> 70
<212> DNA
<213> Artificial Sequence

40 <220>
<223> Description of Artificial Sequence: primer

45 <400> 8
ggagtacagc cactgcgctt ggaccatcgt acgcgtggag atcctgcgca acttctactt 60
catcaaccgc 70

50 <210> 9
<211> 70
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer

55 <400> 9
caccacactg gactagtgga tccttatcag ttgcgcaggt agccggtcag gcggttgatg 60
aagtagaagt 70

60 <210> 10
<211> 70

5 <212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer

10 <400> 10
aggcgcagtg gctgtactcc ttggccttca ggtagtgcag gatgcggcca tagtagcgct 60
tcaggtgcag 70

15 <210> 11
<211> 70
<212> DNA
<213> Artificial Sequence

20 <220>
<223> Description of Artificial Sequence: primer

25 <400> 11
ctccttctcc agcttctcct ccagcacggt cttcaggtgg ttgatctggt ggtacacgct 60
ggccagcagg 70

30 <210> 12
<211> 70
<212> DNA
<213> Artificial Sequence

35 <220>
<223> Description of Artificial Sequence: primer

40 <400> 12
gagctggagt cctggcggaa gatggcgaag atgttctgca gcatctcgta gatggtcaga 60
gcggcgctct 70

45 <210> 13
<211> 70
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer

50 <400> 13
cctcggggat gtcgaagttc atcctgtcct tcaggcagta ctccaggcgc ccgttcagct 60
gccacaggag 70

55 <210> 14
<211> 70
<212> DNA
<213> Artificial Sequence

60 <220>
<223> Description of Artificial Sequence: primer

5
 <400> 14
 caggaagccg agcaggttgt agtcatcga tagggccgtg gtgctgaagc acaggagcag 60
 ggcgatctgg 70

10
 <210> 15
 <211> 70
 <212> DNA
 <213> Artificial Sequence

15
 <220>
 <223> Description of Artificial Sequence: primer

20
 <400> 15
 ctgctccaga tgcacctgct cctgtgcttc agcaccacgg ccctatcgat gaagcaccag 60
 caccagcatc 70

25
 <210> 16
 <211> 70
 <212> DNA
 <213> Artificial Sequence

30
 <220>
 <223> Description of Artificial Sequence: primer

35
 <400> 16
 cactgcttac tggcttatcg aaattaatac gactcactat agggagaccc aagctggcta 60
 gcgtttaaac 70

40
 <210> 17
 <211> 70
 <212> DNA
 <213> Artificial Sequence

45
 <220>
 <223> Description of Artificial Sequence: primer

50
 <400> 17
 caggaagccg agcaggttgt agtcatctg ttggtgttga tggttggtgct gatgctggtg 60
 ctggtgcttc 70

55
 <210> 18
 <211> 70
 <212> DNA
 <213> Artificial Sequence

60
 <220>
 <223> Description of Artificial Sequence: primer

60
 <400> 18
 agcagggcga tctggagcag gcacttggtg gtcagtgtgg cgaagcttaa gtttaaaccg 60
 tagccagctt 70

5	<210> 19	
	<211> 40	
	<212> DNA	
	<213> Artificial Sequence	
10	<220>	
	<223> Description of Artificial Sequence: primer	
	<400> 19	
15	ccgtcagatc ctaggctagc ttattgcggt agtttatcac	40
	<210> 20	
	<211> 32	
20	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Description of Artificial Sequence: primer	
25	<400> 20	
	gagctcggta ccaagctttt aagagctgta at	32
30	<210> 21	
	<211> 77	
	<212> DNA	
	<213> Artificial Sequence	
35	<220>	
	<223> Description of Artificial Sequence: primer	
	<400> 21	
40	gctgaacggg cgcttggagt actgcctgaa ggacaggatg aacttcgaca tccccgagga	60
	aatccgccag ctgcagc	77
	<210> 22	
	<211> 35	
45	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Description of Artificial Sequence: primer	
50	<400> 22	
	tctccacgcg tacgatggtc caggcgaggt ggctg	35
55	<210> 23	
	<211> 70	
	<212> DNA	
	<213> Artificial Sequence	
60	<220>	
	<223> Description of Artificial Sequence: primer	

5

```
<400> 23
caccacactg gactagtgga tccttatcag ttgcgcaggt agccggtcag gcggttgatg 60
aagtagaagt                                     70
```

10

```
<210> 24
<211> 31
<212> DNA
<213> Artificial Sequence
```

15

<220>
<223> Description of Artificial Sequence: primer

20

```
<400> 24
catcagcttg ccggtgggtg tgtcctcctt c 31
```

25

```
<210> 25
<211> 31
<212> DNA
<213> Artificial Sequence
```

30

<220>
<223> Description of Artificial Sequence: primer

<400> 25
gaaggaggac aacaccaccg gcaagctgat g 31

35

```
<210> 26
<211> 40
<212> DNA
<213> Artificial Sequence
```

40

<220>
<223> Description of Artificial Sequence: primer

45

```
<400> 26
cacactggac tagtaagctt ttatcagttg cgcaggtagc 40
```

50

```
<210> 27
<211> 47
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Description of Artificial Sequence: primer

55

<400> 27
gaggagttcg aacttccagt gccagcgcct cctgtggcag ctgaacg 47

60

<210>	28
<211>	29
<212>	DNA

5 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer

10 <400> 28
cgcggatcca tatgaccaac aagtgcctg 29

15 <210> 29
<211> 24
<212> DNA
<213> Artificial Sequence

20 <220>
<223> Description of Artificial Sequence: primer

<400> 29
cgcggatcct tatcagttgc gcag 24

25 <210> 30
<211> 33
<212> DNA
<213> Artificial Sequence

30 <220>
<223> Description of Artificial Sequence: primer

35 <400> 30
tttaaactgg atccagccac catgaccaac aag 33

40 <210> 31
<211> 63
<212> DNA
<213> Artificial Sequence

45 <220>
<223> Description of Artificial Sequence: primer

<400> 31
cggccatagt agcgcttcag gtgcagggag ctcatcagct tgccggtggt gttgtcctcc 60
ttc 63

50 <210> 32
<211> 63
<212> DNA
<213> Artificial Sequence

55 <220>
<223> Description of Artificial Sequence: primer

60 <400> 32
gaaggaggac aacaccaccg gcaagctgat gagctccctg cacctgaagc gctactatgg 60
ccg 63

5

<210> 33
<211> 30
<212> DNA

10 <213> Artificial Sequence

 $\langle 220 \rangle$

<223> Description of Artificial Sequence: primer

15

```
<400> 33
ggcgtcctcc ttggtgaaqt tctgcagctg
```

30

<210> 34

<211> 39

<212> DNA

<213> Artificial Sequence

$\langle 220 \rangle$

25 <223> Description of Artificial Sequence: primer

```
<400> 34
atatatccca agctttttatc agttgcgcag gtagccggt
```

39

<210> 35

<211> 30

<212> DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> Description of Artificial Sequence: primer

```
<400> 35
cagctgcaga acttcaccaa ggaggacgcc
```

30

<210> 36

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 36
cgcggatcca gccaccatga ccaacaagtg cctg

34

<210> 37

<211> 89

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

5

<400> 37
aactggatcc agccaccatg accaacaagt gcctgctcca gatcgccctg ctcctgtgct 60
tcagcaccac ggccctagcc cagagctac 89

10

<210> 38
<211> 92
<212> DNA
<213> Artificial Sequence

15

<220>
<223> Description of Artificial Sequence: primer

20

<400> 38
aactggatcc agccaccatg accaacaagt gcctgctcca gatcgccctg ctcctgtgct 60
tcagcaccac ggccctagcc cagatgagct ac 92

005220"0334360